

Understanding Bed Sores: Pathophysiology and Nursing Interventions

Meshari Hussain Burayk Alotaibi¹, Hussain Raddah Saeyed Alrubaei², Abed Radah Saeed Alrobaie³, Mohsin Mutalq Alotibi⁴, Abdulaziz Helal Difallah Al Zaidi⁵, Maher Dughaylib Almutairi⁶, Fayez Safar Alotaibi⁷, Haitham Owedah Aljehani⁸

¹ Nurse - Erada & Mental Health Complex - Makkah AlMukarramah

² Nurse - Branch of the Ministry of Health in Taif - Makkah AlMukarramah-Taif

³ Nurse - Complex Eradah and Mental Health - Makkah AlMukarramah-Taif

⁴ Nurse - Complex Eradah and Mental Health - Makkah AlMukarramah-Taif

⁵ Nurse - Eradah Complex and Mental Health - Taif - Makkah AlMukarramah

⁶ Nurse - Ministry of Health Branch - Makkah AlMukarramah

⁷ Nursing Technician - Irada Hospital and Mental Health - Taif ⁸

Nursing Specialist - Alamal Hospital - Jeddah

Abstract:

Bed sores, also known as pressure ulcers or decubitus ulcers, are localized injuries to the skin and underlying tissue that occur due to prolonged pressure, most commonly over bony prominences. The pathophysiology involves a complex interplay of factors, including pressure intensity, duration, and tissue tolerance. When pressure is applied to a specific area, it can constrict blood vessels, leading to ischemia and cell death if not relieved. Other contributing factors include moisture, friction, shear forces, and the patient's overall health status, such as nutrition and mobility. Understanding these underlying mechanisms is crucial for effective prevention and management of bed sores. Nursing interventions play a critical role in preventing bed sores and managing existing ones. Key strategies include regular repositioning of patients to relieve pressure, implementing skin assessments to identify early signs of tissue damage, and maintaining skin hygiene and moisture balance. Nutritional support is also vital, as adequate protein and caloric intake promote healing and tissue integrity. Additionally, utilizing appropriate support surfaces, such as special mattresses and cushions, can help distribute pressure more evenly. Education for both nursing staff and caregivers about the risk factors and signs of pressure ulcers is essential in creating a proactive care plan tailored to individual patient needs.

Keywords: Bed Sores, Pressure Ulcers, Pathophysiology, Nursing Interventions, Ischemia, Tissue Damage, Risk Factors, Repositioning, Skin Assessment, Nutritional Support, Support Surfaces, Care Plan

Introduction:

Bed sores, also known as pressure ulcers or decubitus ulcers, represent a significant health concern, particularly among individuals with limited mobility such as the elderly, hospitalized patients, and those suffering from chronic illnesses. These localized injuries to the skin and underlying tissue are often the result of prolonged pressure on the skin, compounded by factors such as friction,

moisture, and shear forces. Understanding the complex pathophysiology of bed sores is essential for healthcare professionals, particularly nurses, who play a pivotal role in the prevention and management of these painful and often debilitating wounds [1].

The pathophysiology of bed sores is multifaceted. Prolonged pressure on bony prominences—such as the sacrum, heels, and elbows—can compromise blood flow to the affected tissues, resulting in ischemia, inflammation, and tissue necrosis. The pressure often exceeds the capillary perfusion pressure, leading to a cascade of cellular events. Initially, the skin may exhibit non-blanchable erythema, which indicates the onset of dermal injury. If pressure is not alleviated, the injury may progress to more severe forms involving the epidermis, dermis, and even deeper tissues, culminating in full-thickness ulcers. Furthermore, factors such as moisture from perspiration or incontinence, friction from movement, and impaired sensory perception due to neurological conditions exacerbate the risk of ulcer development [2].

Additionally, systemic factors, including malnutrition, anemia, and comorbid conditions like diabetes and vascular disease, weaken the skin's structural integrity and hinder the healing process. Understanding these lagging processes is essential for nursing staff, as they can adopt a holistic approach to patient care that addresses both local and systemic issues [3].

Prevention and management of bed sores require a multidisciplinary approach, but nursing interventions are foundational. Evidence-based strategies include conducting thorough skin assessments, implementing appropriate repositioning schedules, utilizing specialized support surfaces, and emphasizing nutrition. Regular skin assessments allow nurses to identify at-risk patients and the early signs of pressure injury, enabling timely interventions. Developing an individualized repositioning protocol can significantly reduce the duration of pressure on vulnerable areas, while the use of pressure-relieving devices—such as foam mattresses or cushions—provides essential support for the skin [4].

Nutritional assessment is equally critical, as adequate protein and caloric intake contribute to wound healing and skin integrity. Comprehensive education for patients and caregivers on the importance of pressure ulcer prevention also empowers them to take an active role in care. Furthermore, adherence to protocols for managing moisture, friction, and shear forces is imperative in clinical practice [5].

Pathophysiology of Bed Sores:

Bed sores, also known as pressure ulcers or decubitus ulcers, represent a significant health concern, particularly among patients with limited mobility. These injuries to the skin and underlying tissue

occur when constant pressure is applied to specific areas of the body, leading to localized damage. The pathophysiology of bed sores is complex and multifactorial, involving an interplay between pressure, friction, shear forces, and various intrinsic and extrinsic factors that influence a patient's susceptibility to developing these injuries. Understanding the pathophysiological mechanisms guiding the onset of bed sores is vital for healthcare professionals, caregivers, and patients alike to implement preventative strategies and effective management [6].

To understand the pathophysiology of bed sores, one must first appreciate the anatomy of the skin. The skin consists of three primary layers: the epidermis (outer layer), the dermis (middle layer), and subcutaneous tissue (innermost layer). Underneath these layers are muscles, fascia, and bones. Bed sores typically occur over bony prominences where the skin is in contact with hard surfaces, such as the sacrum, heels, elbows, and hips [7].

Bony prominences are particularly vulnerable due to their limited cushioning by subcutaneous fat and their proximity to bones, which makes the overlying skin susceptible to pressure-induced damage. The blood supply to the skin is critical for maintaining its integrity and healing capabilities. When tissue perfusion is compromised due to sustained pressure or shearing forces, ischemic injury occurs, leading to cell death and ulceration [7].

Mechanisms of Pressure-Induced Ischemia Pressure

The primary mechanism behind the development of bed sores is pressure. Prolonged pressure on the skin compresses the microcirculation, which impairs blood flow and results in ischemia. Sustained ischemia leads to tissue hypoxia, deprivation of essential nutrients, and the accumulation of metabolic waste products like carbon dioxide and lactic acid. The threshold for pressure that induces ulceration varies among individuals but is commonly accepted to range between 32 mmHg to 50 mmHg, depending on the duration of the pressure applied. When pressure is relieved, there can be reperfusion injury, where the resumption of blood flow leads to oxidative stress and further tissue damage [7].

Friction and Shear Forces

In addition to pressure, friction and shear forces also play crucial roles in the pathophysiology of bed sores. Friction occurs when two surfaces rub against each other; for example, when a patient shifts in bed or is repositioned. This action can result in the abrasion of the skin, stripping away the epidermal layer and exposing the underlying dermis [8].

Shear forces occur when the skin moves in one direction while the underlying structures, such as muscles and bones, remain stationary. This displacement can stretch and tear blood vessels, further compromising blood supply to the region. The combined effects of pressure, friction, and shear forces create an environment conducive to the development of tissue injury [8].

Intrinsic Factors Impacting Bed Sore Development

Several intrinsic factors influence an individual's risk of developing bed sores. These include age, underlying health conditions, nutritional status, and skin moisture.

Age

Older adults are particularly at risk for bed sores due to several reasons. Aging skin experiences changes in elasticity, thickness, and moisture levels, making it more susceptible to damage.

Furthermore, the elderly often have comorbid conditions that can impair mobility and compromise blood flow to the skin, increasing their likelihood of ulceration [9].

Health Conditions

Certain health conditions—such as diabetes mellitus, peripheral vascular disease, and spinal cord injuries—complicate the pathophysiology of pressure ulcers. Diabetes, for example, contributes to poor circulation and nerve damage, making patients unaware of discomfort that would otherwise prompt repositioning [9].

Nutritional Status

Malnutrition can diminish the body's ability to repair tissues and maintain skin integrity. Critical nutrients, such as proteins, vitamins A and C, and zinc play essential roles in collagen synthesis, immune function, and wound healing. A deficiency in these nutrients can substantially increase the risk of pressure ulcer formation [9].

Skin Moisture

Moisture from urine, feces, sweat, or wound drainage can result in skin maceration, weakening the skin barrier and making it more prone to breakdown under pressure. Managing humidity levels in the skin is thus an essential component of prevention.

Extrinsic Factors in Bed Sore Development

Extrinsic factors encompass environmental influences and practices that can contribute to the formation of bed sores. These include the quality of the mattress and bedding, mobility aids, and caregiving practices [10].

Surfaces

The type of mattress and surface on which a patient rests significantly impacts the risk of bed sore development. Hospital-grade mattresses and specialized pressure-relieving devices help distribute body weight and reduce pressure on vulnerable areas. Conversely, inadequate or worn-out bedding can exacerbate the risk [10].

Caregiver Practices

The frequency and methods of repositioning patients also play a pivotal role in preventing bed sores. Regular repositioning—at least every two hours for immobile patients—can help mitigate prolonged pressure. Effective caregiving practices should also involve regular skin assessments and the use of moisture barriers to protect against skin breakdown [11].

Clinical Stages of Bed Sores

Bed sores are classified into stages based on the depth of tissue damage. The National Pressure Injury Advisory Panel (NPIAP) defines the stages as follows:

1. **Stage I:** Non-blanchable erythema is present, indicating skin damage without skin loss. The area may feel warmer, cooler, firmer, or softer than adjacent tissue.
2. **Stage II:** Partial-thickness loss of skin occurs, presenting as a shallow open ulcer with a red or pink wound bed. Blisters may also be present.
3. **Stage III:** Full-thickness tissue loss is apparent, extending through the dermis and into the subcutaneous tissue. Slough may be present but does not obscure the depth of the tissue loss.
4. **Stage IV:** Full-thickness tissue loss extends into muscle, bone, or supporting structures. This stage includes significant damage and often involves necrosis.
5. **Unstageable:** This term is applied when the base of the ulcer is covered by slough or eschar, preventing accurate assessment of the depth [11].

Risk Factors for Bed Sores:

Bed sores, also known as pressure ulcers or decubitus ulcers, are localized injuries to the skin and underlying tissue that occur when there is prolonged pressure on the skin. They are a significant health concern, especially for individuals with limited mobility, and can lead to severe medical complications, including infections and chronic pain. Understanding the risk factors associated with bed sores is critical for prevention and management, particularly in vulnerable populations such as the elderly, individuals with disabilities, and patients in healthcare settings [12].

1. Immobilization and Limited Mobility

The most critical risk factor for the development of bed sores is immobility. Prolonged periods in the same position can compress blood vessels, leading to decreased blood flow to the skin and surrounding tissues. Patients who are bedridden, wheelchair-bound, or have limited ability to move due to surgery, illness, or neurological conditions are at a heightened risk. For instance, individuals after a hip fracture often experience restricted mobility and may remain in one position for extended periods, increasing their susceptibility to pressure injuries [13].

2. Age

Age is another significant factor influencing the risk of bed sores. The skin of elderly individuals tends to be thinner, less elastic, and more fragile than that of younger adults. Additionally, older adults often have multiple comorbidities that contribute to overall frailty, affecting their mobility and hydration status. Age-related changes in circulation can impede the body's ability to deliver

oxygen and nutrients to tissues, further exacerbating the risk of skin breakdown. As such, geriatric patients in nursing homes or receiving home care are particularly vulnerable to developing pressure ulcers [13].

3. Nutritional Deficiency

Proper nutrition plays a vital role in maintaining skin integrity and promoting healing. Malnutrition can weaken the skin and reduce its resistance to pressure injuries. Key nutrients such as proteins, vitamins (particularly vitamins A and C), and minerals (including zinc) are essential for skin health and wound healing. A diet lacking in these nutrients may result from various factors, such as gastrointestinal disorders, difficulty swallowing, or a lack of access to appropriate food. Furthermore, dehydration can lead to dry skin, which is more prone to breaking down under pressure, making hydration another critical component of nutrition in preventing bed sores [13].

4. Moisture and Skin Care

Excess moisture from perspiration, incontinence, or wound drainage can significantly increase the risk of bed sores. Moist skin is more susceptible to friction and shear forces, which can compromise the skin barrier and lead to ulceration. Individuals with incontinence may have prolonged exposure to moisture that irritates the skin and weakens its integrity. Therefore, maintaining a clean and dry environment for patients, along with appropriate skin care products, is essential in minimizing risk factors related to moisture [14].

5. Circulatory Issues

Compromised circulation can impair oxygen and nutrient delivery to tissues, increasing the likelihood of pressure ulcer formation. Conditions such as diabetes, peripheral vascular disease, and congestive heart failure can lead to poor circulation, particularly in the extremities. This reduced blood flow can contribute to a higher susceptibility to tissue ischemia when pressure is applied. Identifying patients with such circulatory issues can assist healthcare providers in implementing more vigilant preventive measures against bed sores [14].

6. Sensory Perception Impairment

Individuals with impaired sensory perception often cannot feel pain or discomfort, which are crucial indicators for repositioning when pressure builds. Neurological conditions, spinal cord injuries, and certain mental health disorders can diminish an individual's awareness of pressure on their skin. This often results in extended periods of immobility as the individual is unable to recognize the need to shift their weight, consequently increasing their risk of developing bed sores. Regularly assessing patients' sensory perception and providing consistent reminders for repositioning can help mitigate this risk [14].

7. Friction and Shear Forces

Friction and shear forces occur when skin is dragged across a surface or slides down when an individual is moved or repositioned. This can result in skin tearing or breaking down, particularly

in vulnerable areas. For example, transferring a patient from a bed to a chair without adequate support can lead to friction injuries. To minimize the risk associated with these forces, healthcare providers should utilize proper lifting techniques and supportive equipment, such as slide sheets or specialized mattresses, which can reduce friction and distribute pressure more evenly [15].

8. Comorbid Conditions

Individuals with multiple health conditions may face an increased risk of developing bed sores due to a combination of factors already discussed. Conditions such as diabetes, cardiovascular diseases, and obesity can complicate healing and impair mobility. Comorbidities can also impact a patient's overall health status, leading to prolonged hospitalization and increased dependency on caregivers. Understanding the interplay between these conditions and pressure ulcer risk can guide personalized preventive interventions [15].

Assessment and Diagnosis:

Pressure sores, also known as pressure ulcers or decubitus ulcers, are localized injuries to the skin and underlying tissue that occur as a result of prolonged pressure, often in combination with shear and friction. These injuries are a significant concern within healthcare settings, particularly for individuals with limited mobility, and they can have severe consequences, including infection and systemic illness. Understanding the evaluation and diagnosis of pressure sores is crucial for healthcare professionals to provide optimal care, improve patient outcomes, and reduce healthcare costs associated with these preventable conditions [16].

Understanding Pressure Sores

Pressure sores commonly develop over bony prominences, such as the sacrum, heels, elbows, and hips, where the pressure from body weight obstructs blood flow to the area. Prolonged pressure can cause the skin and underlying tissue to break down, leading to ulcer formation. Various factors contribute to the development of pressure sores, including immobility, moisture, malnutrition, and reduced sensation [16].

The classification of pressure sores is generally based on depth of tissue loss, as outlined by the National Pressure Injury Advisory Panel (NPIAP). There are four stages of pressure ulcers:

- **Stage I:** Non-blanchable erythema of intact skin. The area may be painful, firm, soft, warm, or cool compared to surrounding tissue.
- **Stage II:** Partial thickness loss of skin with exposed dermis. This stage may present as a blister or shallow open sore.
- **Stage III:** Full thickness loss of skin, extending into the subcutaneous tissue but not through the underlying fascia. The ulcer may present as a deep crater.
- **Stage IV:** Full thickness tissue loss with exposed bone, tendon, or muscle. Slough or eschar may be present on some parts of the wound bed [17].

In addition to these stages, pressure injuries may be categorized as unstageable when the base of the ulcer is covered by slough or eschar, making it impossible to assess the depth accurately.

Risk Factors

Identifying individuals at risk for pressure sore development is a crucial element in the evaluation process. Several intrinsic and extrinsic factors contribute to the risk, including:

1. Intrinsic Factors:

- **Immobility:** The inability to change positions can impede blood flow and increase pressure on specific body areas.
- **Sensory Perception Impairment:** Conditions such as spinal cord injuries, stroke, or neuropathies can limit a person's ability to feel discomfort or pain, making them unaware of prolonged pressure.
- **Nutritional Status:** Malnutrition and dehydration can compromise skin integrity and healing capacity, increasing susceptibility to ulcer formation.
- **Age:** The elderly are more at risk due to thinning skin, decreased blood circulation, and other age-related health conditions [18].

2. Extrinsic Factors:

- **Moisture:** Excessive moisture from incontinence or perspiration can weaken skin, making it prone to breakdown.
- **Friction and Shear:** Sliding down in bed or being dragged across surfaces can damage top layers of skin and exacerbate pressure injury development [18].

Evaluation Process

The evaluation of pressure sores involves a comprehensive assessment that includes taking a detailed medical history, performing a physical examination, and utilizing validated assessment tools.

1. **Medical History:** A thorough history should be gathered, including risk factors, past occurrences of pressure sores, comorbid conditions, current medications, and nutritional status. Assessment scales, like the Braden Scale or Norton Scale, can assist in determining the risk level for pressure sores [19].
2. **Physical Examination:** The examination should focus on identifying existing sores and assessing their stage, size, location, drainage, pain level, and surrounding tissue condition. Observation of factors such as temperature, color, and texture of the skin is crucial [19].
3. **Wound Assessment:** Recording the dimensions of the ulcer, including length, width, and depth, is essential for monitoring progress and response to treatment. Additionally,

assessing the wound bed and surrounding tissue for signs of erythema, edema, or signs of infection (such as purulent drainage or foul odor) can help in deciding further management strategies.

4. **Diagnostic Tests:** In cases of suspected infection, cultures may be performed to identify pathogenic organisms. Blood tests may also be ordered to evaluate for systemic involvement, rule out underlying conditions, or assess nutritional status [19].

Diagnosis

The diagnosis of pressure sores is primarily clinical and relies on the findings from the evaluation process. The healthcare provider must accurately stage the ulcer based on standardized criteria to guide treatment decisions. Documenting the stage, size, and overall condition of the wound is essential for future comparisons and evaluating the effectiveness of interventions.

Furthermore, diagnosing pressure sores necessitates continuous reassessment. Regular evaluations should be scheduled to monitor for changes in clinical status, especially as healing progresses or if new pressure sores develop [20].

Preventive Nursing Interventions:

Bed sores, also known as pressure ulcers or decubitus ulcers, represent a significant health concern, particularly for individuals with limited mobility, such as those who are bedridden, elderly, or suffering from debilitating medical conditions. These injurious skin lesions occur due to prolonged pressure on the skin, typically in areas where the bones are close to the skin's surface, such as the heels, sacrum, and elbows. The implications of bed sores extend beyond physical discomfort; they can lead to serious infections, prolonged hospital stays, decreased quality of life, and increased healthcare costs. Therefore, preventive nursing interventions play a crucial role in mitigating the risk of developing bed sores [21].

Identifying individuals at risk is a critical step in preventing bed sores. Key risk factors include immobility, poor nutrition, advanced age, moisture, and chronic medical conditions like diabetes and vascular diseases. Other contributing factors can include skin friction and shear, incontinence, and insufficient health care resources. Nurses must assess these factors through regular evaluations, using tools such as the Braden Scale, which measures a patient's risk for pressure ulcers by evaluating sensory perception, moisture, activity, mobility, nutrition, and friction/shear [21].

Regular skin assessment is essential in preventing bed sores. Nurses should conduct thorough skin checks at least once a day or more frequently for high-risk patients. The assessment should focus on bony prominences and areas most prone to pressure ulcers, inspecting for any redness, warmth, swelling, or breakdown of the skin. Implementing a standardized skin assessment protocol can ensure consistency and thoroughness in patient evaluations [22].

Additionally, continuous monitoring of patients allows for timely interventions if any signs of potential pressure ulcers are detected. It helps in understanding how a patient's condition may

change over time and facilitates ongoing evaluation of the effectiveness of preventive measures implemented [22].

Nutrition plays a vital role in skin health and wound healing. Malnutrition can compromise the integrity of the skin and make individuals more susceptible to pressure ulcers. Nurses should assess patients' nutritional status and collaborate with dietitians to devise a meal plan that meets their caloric and protein needs. Nutritional supplementation may be necessary for patients who are unable to consume adequate nutrients, with special attention to vitamins and minerals crucial for skin integrity, such as Vitamin C, Zinc, and protein-rich foods [23].

A consistent and proactive skin care regimen is paramount in preventing bed sores. This regimen may include keeping the skin clean and dry, using gentle, non-irritating cleansers, and ensuring adequate hydration. Frequent changes of incontinence pads can help reduce moisture-related skin irritation. Moreover, emollients or barrier creams can be applied to protect the skin from breakdown, especially in high-risk areas [23].

Regular repositioning is one of the most effective strategies to prevent pressure ulcers. Nurses should establish a turning schedule, repositioning patients at least every two hours when they are in bed, and more frequently when they are sitting. Utilizing specialized mattresses or cushions designed to reduce pressure can also be beneficial. These devices can help distribute weight evenly and relieve pressure on vulnerable areas [24].

Encouraging mobility, when possible, is another important preventive measure. Patients should be assisted in moving, shifting, or even standing up as much as their condition allows. For those who cannot independently change positions, utilizing transfer aids or mechanical devices to facilitate movement can help reduce prolonged pressure on specific areas of the body.

Education is key in empowering patients, families, and caregivers to understand the importance of pressure ulcer prevention. Nurses should provide comprehensive guidance on the significance of repositioning, nutrition, and skin care practices. Additionally, collaboration with the entire healthcare team—including physicians, dietitians, and physiotherapists—is essential in developing a cohesive plan tailored to the individual patient's needs [25].

Management of Existing Pressure Ulcers:

Pressure ulcers, also known as bedsores or decubitus ulcers, are localized injuries to the skin and underlying tissue that occur as a result of prolonged pressure, often over bony prominences. They are most commonly found in individuals with limited mobility, including patients in hospitals, nursing homes, and those at home who are immobile due to illness or disability. The management of existing pressure ulcers is critical not only to promote healing but also to prevent complications such as infections, increased morbidity, and prolonged hospital stays [26].

Understanding Pressure Ulcers

Before delving into management strategies, it is important to understand the classification of pressure ulcers. The National Pressure Injury Advisory Panel (NPIAP) categorizes pressure ulcers into four stages:

1. **Stage 1:** Intact skin with a localized area of non-blanchable erythema.
2. **Stage 2:** Partial-thickness loss of skin with exposed dermis, presenting as a wound bed that is pink or red and moist.
3. **Stage 3:** Full-thickness tissue loss, where subcutaneous fat may be visible, but bone, tendon, or muscle are not exposed.
4. **Stage 4:** Full-thickness tissue loss with exposed bone, tendon, or muscle, often accompanied by necrosis or damage to surrounding tissues [27].

Preventing the progression of existing ulcers and fostering healing is vital. The management protocol for existing ulcers generally includes thorough assessment, adequate nutrition, wound care, pain management, and comprehensive rehabilitation [28].

Assessment of Pressure Ulcers

A thorough assessment of pressure ulcers is pivotal for devising an effective management plan. Clinicians should first identify the stage of the ulcer, documenting its size, depth, location, and wound characteristics such as exudate levels and condition of the surrounding skin.

Several standardized tools can assist healthcare providers in assessing and documenting pressure ulcers. The Braden Scale is one example, which evaluates risk factors such as sensory perception, moisture, activity, mobility, nutrition, and friction/shear to predict ulcer risk. Frequent reassessments are essential, particularly after notable changes in the patient's condition or after interventions implemented for management [29].

Nutritional Considerations

Nutrition plays a crucial role in wound healing. Malnutrition can hinder the body's ability to repair tissue, and individuals with pressure ulcers may have increased caloric and protein needs. A diet rich in proteins, vitamins, and minerals is essential. Key nutrients include:

- **Protein:** Vital for tissue repair; sources include lean meats, dairy, and legumes [30].
- **Vitamin C:** Critical for collagen synthesis; foods such as oranges, strawberries, and green vegetables are beneficial.
- **Zinc:** Plays a role in immune function and collagen synthesis; sources include meat, shellfish, and nuts.
- **Vitamin A:** Important for skin health, found in foods like carrots, sweet potatoes, and greens.

Healthcare providers should consider consulting a dietitian to develop an individualized nutrition plan [30].

Wound Care Strategies

Effective wound care is a cornerstone of pressure ulcer management. Several approaches aim to maintain a moist wound environment, which is conducive to healing:

1. **Cleansing:** The wound should be cleaned gently with saline or a non-toxic wound cleanser. Harsh antiseptics should be avoided as they can further damage tissue [31].
2. **Dressing Choices:** Various dressings are available depending on the ulcer's stage and characteristics. Hydrocolloid, foam, alginate, and transparent film dressings all serve different purposes:
 - **Hydrocolloid dressings:** Provide a moist environment and are suitable for stage 2 ulcers.
 - **Foam dressings:** Absorb exudate and are appropriate for stage 3 ulcers.
 - **Alginate dressings:** Ideal for highly exudative wounds, allowing for moisture balance [31].
3. **Debridement:** For ulcers with necrotic tissue, debridement is important to remove dead tissue and promote healing. Methods may include surgical debridement, enzymatic debridement, or autolytic debridement.
4. **Infection Control:** Signs of infection such as increased redness, warmth, swelling, and purulent drainage necessitate culture and appropriate antimicrobial treatment. Topical or systemic antibiotics may be required, depending on the severity of the infection [31].

Pain Management

Patients with pressure ulcers often experience pain, which can be detrimental to their overall wellbeing and impair healing. Effective pain management strategies should be tailored to the individual and may involve pharmacologic options such as analgesics, anti-inflammatory drugs, and topical anesthetics. Additionally, non-pharmacologic measures like repositioning, relaxation techniques, and distraction methods can be beneficial [32].

The management of pressure ulcers requires a collaborative approach involving a multidisciplinary team. In addition to nurses, physicians, and dietitians, wound care specialists, physical and occupational therapists, and psychologists may also play essential roles. Physical therapists can assist in improving mobility and repositioning techniques to alleviate pressure on sensitive areas. Occupational therapists can help implement adaptive equipment that encourages independence while minimizing pressure risks [33].

Patient and Caregiver Education

Education is an essential component of managing pressure ulcers. Patients and caregivers should be informed about the importance of routine skin assessment and the early signs of pressure ulcer development. Educational resources should also emphasize the need for regular repositioning and techniques to relieve pressure, as well as the critical role of nutrition in healing [33].

Role of Nutrition in Pressure Ulcer Prevention and Healing:

Pressure ulcers, also known as bedsores or decubitus ulcers, are localized injuries to the skin and underlying tissue that occur due to prolonged pressure, often in individuals with limited mobility. These injuries can range from mild skin discoloration to severe tissue damage, including deep wounds that can become infected. The incidence of pressure ulcers is particularly high in elderly populations, patients with spinal cord injuries, and individuals with chronic illnesses. While the prevention and management of pressure ulcers primarily emphasize repositioning, skin care, and wound management, an often underappreciated factor in both prevention and healing is nutrition [34].

Pressure ulcers occur when pressure exceeds the capillary perfusion pressure, thereby restricting blood flow to the tissues. This lack of blood flow can lead to ischemia, cell death, and a subsequent breakdown of skin integrity. Key risk factors include immobility, moisture, friction, and an individual's overall health status. With nearly 2.5 million individuals affected by pressure ulcers annually in the United States alone, this condition represents not only a significant health challenge but also a considerable economic burden, leading to increased healthcare costs and prolonged hospital stays [35].

Nutritional Requirements for Skin Integrity

Skin is the body's largest organ, and its integrity is largely dependent on the presence of vital nutrients. Various nutrients play distinct roles in maintaining skin health, preventing the occurrence of pressure ulcers, and facilitating their healing process [35].

1. Proteins

Proteins are fundamental for cell growth and repair. They are vital in the wound healing process, as they contribute to tissue regeneration. The amino acids derived from proteins are the building blocks for new tissue and play a crucial role in collagen synthesis. Collagen is an essential protein that helps maintain skin structure and strength. Inadequate protein intake can substantially slow down the healing of pressure ulcers, making the recovery process longer and more complicated. Thus, ensuring a sufficient intake of protein – particularly high-quality sources such as lean meats, dairy, legumes, and nuts – is critical in preventing ulcer formation and promoting healing [36].

2. Calories

For effective healing, the body requires an adequate caloric intake to provide energy for cellular processes, including repair and regeneration. Individuals with existing pressure ulcers often need additional calories to support the increased metabolic demand associated with healing. An insufficient caloric intake may lead to weight loss, muscle wasting, and

decreased strength, all of which can hinder mobility and increase the risk of further skin breakdown [37].

3. Vitamins and Minerals

Vitamins and minerals are essential co-factors in numerous biochemical processes involved in skin health and wound healing. Key vitamins include:

- **Vitamin A:** Vital for cell proliferation and differentiation, Vitamin A helps in maintaining the integrity of the skin and mucous membranes. It also assists in the inflammatory response, which is crucial during the early stages of wound healing.
- **Vitamin C:** An antioxidant that plays a crucial role in collagen synthesis and also helps to reduce oxidative stress, thus aiding in wound healing. Deficiency in Vitamin C can lead to impaired healing and increased susceptibility to infections.
- **Vitamin E:** Another antioxidant that helps protect cell membranes from oxidative damage and plays a role in the inflammatory response.
- **Zinc:** Essential for protein synthesis and cell proliferation, zinc deficiencies can lead to delayed wound healing and increased risk of infections, compounding the issues associated with pressure ulcers [38].

4. Hydration

Proper hydration is another critical component of nutrition that supports skin integrity. Dehydration can lead to dry skin, making it more susceptible to breakdown. Adequate fluid intake is necessary for maintaining skin turgor and elasticity. Additionally, hydration aids in the overall metabolic processes required for healing [39].

5. Fatty

Acids

Essential fatty acids, particularly omega-3 and omega-6 fatty acids, play roles in antiinflammatory responses and cellular repair. These fatty acids can help reduce the inflammation associated with pressure ulcers and support overall skin health [40].

Nutritional Assessment and Interventions

Given the significant role of nutrition in pressure ulcer prevention and healing, healthcare providers must conduct thorough nutritional assessments for at-risk patients. This assessment should evaluate individual dietary intake, nutritional status, and any barriers to proper nutrition [41].

If nutritional deficiencies are identified, targeted interventions should include:

- **Nutritional Counseling:** Providing education on the importance of balanced nutrition and how it relates to skin health.

- **Dietary Modifications:** Tailoring meal plans to ensure adequate calorie, protein, vitamin, and mineral intake based on individual needs.
- **Supplements:** In some cases, dietary supplements may be necessary to provide essential nutrients that are insufficient in the diet, particularly for patients with increased nutritional needs or those who have difficulty eating [42].

Education and Training for Healthcare Providers:

Pressure ulcers, also known as pressure sores or bedsore, are localized injuries to the skin and underlying tissue, usually over bony prominences, due to prolonged pressure, shear, and friction. These injuries not only pose a significant challenge to patient care but also represent a considerable burden to healthcare systems worldwide. The pathophysiology of pressure ulcers is complex and involves multiple factors including ischemia, inflammation, and tissue breakdown. Comprehensive education and training for healthcare providers are vital to not only prevent these injuries but also to enhance patient outcomes [43].

To effectively combat pressure ulcers, healthcare providers must understand the underlying pathophysiology. Pressure ulcers result from a combination of mechanical forces—pressure, shear, and friction—that compromise blood flow to the skin and underlying tissues. Sustained pressure, particularly over areas with minimal soft tissue coverage, such as the sacrum, heels, and elbows, leads to ischemia; a critical reduction in blood flow deprives tissues of oxygen and nutrients, resulting in cellular damage and death [44].

Moreover, shear forces occur when the skin is moved in one direction while the underlying tissues are stationary, often resulting in vascular damage. Friction, while not as significant a contributor, can exacerbate the damage caused by pressure and shear, especially in patients with compromised skin integrity. In addition to these mechanical factors, systemic conditions such as nutritional deficits, moisture, and infection can further complicate the situation. Understanding these multifactorial aspects is imperative for healthcare providers' ability to implement effective risk assessment and management strategies [45].

Effective education and training for healthcare providers are essential for multiple reasons. Firstly, knowledge of the pathophysiology of pressure ulcers equips providers to identify at-risk patients more accurately. Early identification of vulnerable patients—such as those with limited mobility, chronic illness, or malnutrition—enables timely interventions that can prevent ulcer formation [46].

Secondly, comprehensive training ensures that healthcare providers are familiar with risk assessment tools and classification systems. Leveraging standardized tools, such as the Braden Scale, helps facilitate early detection and stratification of risk levels, allowing for tailored management plans. Additionally, education enhances communication among healthcare teams, improving collaborative efforts to optimize patient care [47].

Furthermore, ongoing training associated with the latest evidence-based practices fosters a culture of continuous improvement in patient safety. Regular workshops, seminars, and continuing education programs can keep healthcare providers abreast of advancements in the field, including new materials for wound care, innovative interventions, and the role of technology in monitoring at-risk patients [48].

Current Guidelines and Best Practices

Several organizations provide guidelines designed to reduce the incidence of pressure ulcers, emphasizing the importance of a comprehensive approach that includes risk assessment, prevention strategies, and effective management. The National Pressure Injury Advisory Panel (NPIAP) and The Agency for Healthcare Research and Quality (AHRQ) recommend a multifaceted strategy that includes:

1. **Risk Assessment:** Utilizing validated tools like the Braden Scale to assess patient risk upon admission and with regular intervals throughout care [49].
2. **Skin Assessment:** Conducting routine skin assessments to catch early signs of pressure damage.
3. **Repositioning:** Implementing a schedule for turning and repositioning patients to relieve pressure.
4. **Nutrition:** Ensuring proper nutritional support to promote skin integrity and healing.
5. **Moisture Management:** Implementing effective methods to manage moisture, particularly in patients with incontinence.
6. **Education:** Providing targeted education to both staff and patients regarding pressure ulcer prevention techniques.

These guidelines emphasize teamwork, involvement of patients, and caregivers, as well as the importance of documentation in the management of pressure ulcers [49].

Strategies for Effective Training

To ensure that educational initiatives on the pathophysiology of pressure ulcers are effective, healthcare facilities must consider the following strategies:

1. **Curriculum Development:** Training should be comprehensive, incorporating various learning formats such as lectures, interactive workshops, and simulation-based education [50]. Incorporating case studies will help participants to analyze real-life scenarios and interactive discussions can enhance critical thinking [50].
2. **Interdisciplinary Approach:** Incorporate insights from various disciplines, including nursing, nursing assistants, dietitians, and physical therapists. An interdisciplinary module

promotes an understanding of how different roles contribute to pressure ulcer prevention and management, fostering teamwork [51].

3. **Use of Technology:** Healthcare facilities can utilize e-learning platforms that allow flexible education. Simulations and virtual reality can help in demonstrating the impact of pressure and effective repositioning [52].
4. **Regular Updates and Continuous Education:** Continuous education is crucial in keeping up with advancements in pressure ulcer management and care. Regularly scheduled training sessions and access to online courses can ensure that healthcare staff remain informed about new guidelines and innovations [53].
5. **Evaluation and Feedback:** Develop assessments to evaluate the effectiveness of training programs and obtain feedback from participants. This can help in refining curricula to better align with the needs and learning preferences of the staff [54].
6. **Patient and Family Involvement:** Educating patients and their families on the factors that contribute to pressure ulcers and preventive measures builds awareness and empowers them in their care processes. Effective communication is key to engaging patients in their own healthcare [55].

Conclusion:

In conclusion, understanding bed sores, or pressure ulcers, is vital for improving patient outcomes in clinical settings. The pathophysiology of bed sores involves complex physiological processes where sustained pressure leads to tissue ischemia and subsequent injury, exacerbated by factors such as moisture, friction, and poor nutritional status. By recognizing the risk factors associated with ulcer development, healthcare professionals can implement effective preventive strategies, including regular repositioning, diligent skin assessments, and nutritional support.

Nursing interventions play a crucial role in both preventing and managing bed sores. Continuous education and training for healthcare providers are essential to ensure adherence to best practices and maintain awareness of emerging evidence-based approaches. By fostering a culture of vigilance and responsiveness to patient needs, healthcare teams can significantly reduce the incidence of pressure ulcers, enhance healing outcomes for affected individuals, and ultimately improve the quality of care provided in various healthcare environments. Prioritizing the task of understanding and managing bed sores not only contributes to better patient outcomes but also aligns with the broader goals of patient safety and comfort in nursing practice.

References:

1. Kaveeshwar SA, Cornwall J. The current state of diabetes mellitus in India. *Australas Med J.* 2014;7:45–8. doi: 10.4066/AMJ.2013.1979.

2. Reddy NP, Patel K. A mathematical model of flow through the terminal lymphatics. *Med Eng Phys.* 1995;17:134–40. doi: 10.1016/1350-4533(95)91885-k.
3. Cannon BC, Cannon JP. Management of pressure ulcers. *Am J Health Syst Pharm.* 2004;61:1895–905.
4. Bliss MR. Hyperaemia. *J Tissue Viability.* 1998;8:4–13. doi: 10.1016/s0965206x(98)80028-4.
5. Gefen A. Reswick and Rogers pressure-time curve for pressure ulcer risk. Part 2. *Nurs Stand.* 2009;23:40–4. doi: 10.7748/ns2009.07.23.46.40.c7169.
6. Langemo D, Anderson J, Hanson D, Hunter S, Thompson P, Posthauer ME. Nutritional considerations in wound care. *Adv Skin Wound Care.* 2006;19:297–8. doi: 10.1097/00129334-200607000-00007.
7. Vasconez LO, Schneider WJ, Jurkiewicz MJ. Pressure sores. *Curr Prob Surg.* 1977;62:1–62. doi: 10.1016/s0011-3840(77)80007-5.
8. Callam MJ, Ruckley CV, Harper DR, Dale JJ. Chronic ulceration of the leg: Extent of the problem and provision of care. *Br Med J (Clin Res Ed)* 1985;290:1855–6. doi: 10.1136/bmj.290.6485.1855.
9. Jay R. Pressure and shear: Their effects on support surface choice. *Ostomy Wound Manage.* 1995;41:36–8.
10. Scivoletto G, Fuoco U, Morganti B, Cosentino E, Molinari M. Pressure sores and blood and serum dysmetabolism in spinal cord injury patients. *Spinal Cord.* 2004;42:473–6. doi: 10.1038/sj.sc.3101622.
11. Dodd KT, Gross DR. Three-dimensional tissue deformation in subcutaneous tissues overlying bony prominences may help to explain external load transfer to the interstitium. *J Biomech.* 1991;24:11–9. doi: 10.1016/0021-9290(91)90322-e.
12. Krouskop TA, Reddy NP, Spencer WA, Secor JW. Mechanisms of decubitus ulcer formation — An hypothesis. *Med Hypotheses.* 1978;4:37–9. doi: 10.1016/03069877(78)90025-7.
13. Gefen A. Reswick and Rogers pressure-time curve for pressure ulcer risk. Part 1. *Nurs Stand.* 2009;23:64. doi: 10.7748/ns2009.07.23.45.64.c7115.
14. Keast DH, Fraser C. Treatment of chronic skin ulcers in individuals with anemia of chronic disease using recombinant human erythropoietin (EPO): A review of four cases. *Ostomy Wound Manage.* 2004;50:64–70.

15. Leblebici B, Turhan N, Adam M, Akman MN. Clinical and epidemiologic evaluation of pressure ulcers in patients at a university hospital in Turkey. *J Wound Ostomy Continence Nurs.*
16. Zarei E, Madarshahian E, Nikkhah A, Khodakarim S. Incidence of pressure ulcers in intensive care units and direct costs of treatment: Evidence from Iran. *J Tissue Viability.* 2019;28(2):70–4. doi: 10.1016/j.jtv.2019.02.001.
17. Mervis JS, Phillips TJ. Pressure ulcers: Pathophysiology, epidemiology, risk factors, and presentation. *J Am Acad Dermatol.* 2019;81(4):881–90. doi: 10.1016/j.jaad.2018.12.069.
18. Uchida Y, Kitayama S, Mesquita B, Reyes JAS, Morling B. Is perceived emotional support beneficial? Well-being and health in independent and interdependent cultures. *Pers Soc Psychol Bull.* 2008;34(6):741–54. doi: 10.1177/0146167208315157.
19. Gorecki C, Brown JM, Nelson EA, Briggs M, Schoonhoven L, Dealey C, et al. Impact of pressure ulcers on quality of life in older patients: a systematic review. *J Am Geriatr Soc.* 2009;57(7):1175–83. doi: 10.1111/j.1532-5415.2009.02307.x.
20. Davydow DS, Gifford JM, Desai SV, Needham DM, Bienvenu OJ. Posttraumatic stress disorder in general intensive care unit survivors: a systematic review. *Gen Hosp Psychiatry.* 2008;30(5):421–34. doi: 10.1016/j.genhosppsy.2008.05.006.
21. Bielecki M, Bielecki P, Żebrowski P, Misiak B, Lewko J. Operative treatment of pressure ulcers using pedicled flaps. *Prog Health Sci.* 2018;8(2):105–11.
22. Koo M, Sim Y, Kang I. Risk Factors of Medical Device-Related Pressure Ulcer in Intensive Care Units. *J Korean Acad Nurs.* 2019;49(1):36–45. doi: 10.4040/jkan.2019.49.1.36.
23. Kim HS, Sherman DK, Ko D, Taylor SE. Pursuit of comfort and pursuit of harmony: Culture, relationships, and social support seeking. *Pers Soc Psychol Bull.* 2006;32(12):1595–607. doi: 10.1177/0146167206291991.
24. Holloway I, Galvin K. *Qualitative research in nursing and healthcare.* John Wiley & Sons; 2016.
25. Burhans LM, Alligood MR. Quality nursing care in the words of nurses. *J Adv Nurs.* 2010;66(8):1689–97. doi: 10.1111/j.1365-2648.2010.05344.x.
26. Manojlovich M, Antonakos CL, Ronis DL. Intensive care units, communication between nurses and physicians, and patients' outcomes. *Am J Crit Care.* 2009;18(1):21–30. doi: 10.4037/ajcc2009353.
27. Spilsbury K, Nelson A, Cullum N, Iglesias C, Nixon J, Mason S. Pressure ulcers and their treatment and effects on quality of life: hospital inpatient perspectives. *J Adv Nurs.* 2007;57(5):494–504. doi: 10.1111/j.1365-2648.2006.04140.x.

28. Liao X, Ju Y, Liu G, Zhao X, Wang Y, Wang Y. Risk factors for pressure sores in hospitalized acute ischemic stroke patients. *J Stroke Cerebrovasc Dis.* 2019;28(7):2026–30. doi: 10.1016/j.jstrokecerebrovasdis.2019.02.033.
29. Elliott R, McKinley S, Fox V. Quality improvement program to reduce the prevalence of pressure ulcers in an intensive care unit. *Am J Crit Care.* 2008;17(4):328–34.
30. Okhovati S, Esmaeili M, Shariat E. Effect of Intensive Care Unit Nurses' Empowerment Program on Ability in Visual Differential Diagnosis of Pressure Ulcer Classification. *Crit Care Nurs Q.* 2019;42(1):89–95. doi: 10.1097/CNQ.0000000000000242.
31. Akhkand SS, Seidi J, Ebadi A, Gheshlagh RG. Prevalence of pressure ulcer in Iran's intensive care units: A systematic review and a meta-analysis. *Nurs Pract Today.* 2020;7(1):21–9.
32. Sardari M, Esmaeili R, Ravesh NN, Nasiri M. The impact of pressure ulcer training program on nurses' performance. *J Adv Pharm Educ Res.* 2019;9:145–149.
33. Schreier M. Qualitative content analysis. *The SAGE handbook of qualitative data analysis.* 2014. pp. 170–83.
34. Mojaverian T, Kim HS. Interpreting a helping hand: Cultural variation in the effectiveness of solicited and unsolicited social support. *Personality and Social Psychology Bulletin.* 2013;39(1):88–99. doi: 10.1177/0146167212465319.
35. Young ZF, Evans A, Davis J. Nosocomial pressure ulcer prevention: a successful project. *JONA.* 2003;33(7-8):380–3. doi: 10.1097/00005110-200307000-00004.
36. Serrano ML, Mendez MG, Cebollero FC, Rodriguez JL. Risk factors for pressure ulcer development in Intensive Care Units: A systematic review. *Med Intensiva.* 2017;41(6):339–46. doi: 10.1016/j.medin.2016.09.003.
37. Cobb S. Social support as a moderator of life stress. *Psychosom Med.* 1976;38(5):300–14. doi: 10.1097/00006842-197609000-00003.
38. González Sarasúa J, Pérez López S, Álvarez Viejo M, Pérez Basterrechea M, Fernández Rodríguez A, Ferrero Gutiérrez A, et al. Treatment of pressure ulcers with autologous bone marrow nuclear cells in patients with spinal cord injury. *J Spinal Cord Med.* 2011;34:301–7. doi: 10.1179/2045772311Y.0000000010.
39. Al Fadhli A, Alexander G, Kanjoor JR. Versatile use of vacuum-assisted healing in fifty patients. *Indian J Plast Surg.* 2009;42:161–8. doi: 10.4103/0970-0358.59273.
40. Kamata Y, Takahashi Y, Iwamoto M, Matsui K, Murakami Y, Muroi K, et al. Local implantation of autologous mononuclear cells from bone marrow and peripheral blood for

- treatment of ischaemic digits in patients with connective tissue diseases. *Rheumatology (Oxford)* 2007;46:882–4. doi: 10.1093/rheumatology/kel436.
41. Mayfield JA, Reiber GE, Sanders LJ, Janisse D, Pogach LM. Preventive foot care in people with diabetes. *Diabetes Care*. 1998;21:2161–77. doi: 10.2337/diacare.21.12.2161.
 42. Mumcuoglu KY, Lipo M, Ioffe-Uspensky I, Miller J, Galun R. Maggot therapy for gangrene and osteomyelitis. *Harefuah*. 1997;132:323–5.
 43. Sherman RA. Maggot versus conservative debridement therapy for the treatment of pressure ulcers. *Wound Repair Regen*. 2002;10:208–14. doi: 10.1046/j.1524475X.2002.10403.x.
 44. Desneves KJ, Todorovic BE, Cassar A, Crowe TC. Treatment with supplementary arginine, vitamin C and zinc in patients with pressure ulcers: A randomised controlled trial. *Clin Nutr*. 2005;24:979–87. doi: 10.1016/j.clnu.2005.06.011.
 45. Kavros SJ, Liedl DA, Boon AJ, Miller JL, Hobbs JA, Andrews KL. Expedited wound healing with noncontact, low-frequency ultrasound therapy in chronic wounds: A retrospective analysis. *Adv Skin Wound Care*. 2008;21:416–23. doi: 10.1097/01.ASW.0000323546.04734.31.
 46. Ramundo J, Gray M. Enzymatic wound debridement. *J Wound Ostomy Continence Nurs*. 2008;35:273–80. doi: 10.1097/01.WON.0000319125.21854.78.
 47. Webb LX. New techniques in wound management: Vacuum-assisted wound closure. *J Am Acad Orthop Surg*. 2002;10:303–11. doi: 10.5435/00124635-200209000-00002.
 48. Leaper DJ. Silver dressings: Their role in wound management. *Int Wound J*. 2006;3:282–94. doi: 10.1111/j.1742-481X.2006.00265.x.
 49. Fletcher J. The benefits of using hydrocolloids. *Nurs Times*. 2003;99:57.
 50. Moore ZE, Cowman S. Wound cleansing for pressure ulcers. *Cochrane Database Syst Rev*. 2005;3:CD004983. doi: 10.1002/14651858.CD004983.pub2.
 51. Wasiak J, Cleland H, Campbell F. Dressings for superficial and partial thickness burns. *Cochrane Database Syst Rev*. 2008;4:CD002106. doi: 10.1002/14651858.CD002106.pub3.
 52. Graham JS, Schomacker KT, Glatter RD, Briscoe CM, Braue EH, Jr, Squibb KS. Efficacy of laser debridement with autologous split-thickness skin grafting in promoting improved healing of deep cutaneous sulfur mustard burns. *Burns*. 2002;28:719–30. doi: 10.1016/s0305-4179(02)00198-5.
 53. Ramundo J, Gray M. Is ultrasonic mist therapy effective for debriding chronic wounds? *J Wound Ostomy Continence Nurs*. 2008;35:579–83. doi: 10.1097/01.WON.0000341470.41191.51.

Meshari Hussain Burayk Alotaibi¹, Hussain Raddah Saeyed Alrubaei², Abed Radah Saeed Alrobaie³, Mohsin Mutalq Alotibi⁴, Abdulaziz Helal Difallah Al Zaidi⁵, Maher Dughaylib Almutairi⁶, Fayez Safar Alotaibi⁷, Haitham Owedah Aljehani⁸

54. Berlowitz DR, Brandeis GH, Anderson J, Brand HK. Predictors of pressure ulcer healing among long-term care residents. *J Am Geriatr Soc.* 1997;45:30–4. doi: 10.1111/j.15325415.1997.tb00974.x.
55. Molan PC. The evidence supporting the use of honey as a wound dressing. *Int J Low Extrem Wounds.* 2006;5:40–54. doi: 10.1177/1534734605286014.