

The Role of Imaging Modalities on the Diagnosis and Management of Odontogenic Cysts

Mohammed Ahmed Alhudiry¹, Mohammed Ibrahim Fageeh², Abdullah Hussain Alasmari³, Mohammed Ali Ahmed Assiri⁴, Faisal Hadi Abdullah Asiri⁵, Abdullah Mohammed Alsayed⁶, Mohammed Ibrahim Muidh Althui⁷, Nahla Atia Nasser Albarakati⁸, Hasnaa Eshaq Haroun Hawsawi⁹, Hashem Alawi Melebari¹⁰, Kholood Harisi¹¹, Ohood Abdaraboh Alhabashi¹², Nasir Abduh Mana Al Ahmari¹³, Amal Saleh Alhiani¹⁴, Wafaa Mohammed Alomari¹⁵

¹Dentist, Asseer central hospital

²Dentist, Asseer central hospital

³Muhayil Specialized dental Center, General dentist

⁴Muhayil Specialized dental Center, General dentist

⁵Muhayil Specialized Dental Center, Dental technician

⁶Muhayil Specialist Dental Center, Dental assistant

⁷Muhayil Specialized Dental Center, Dental Assistant Technician

⁸Dental Technician, Al- aziziyah urgent care Makkah

⁹Dental assistant technician, Alhindawiya primary healthcare Makkah

¹⁰DENTAL ASISTANT, Executive management of supply chains in the Makkah cluster

¹¹Dental assistant, King Abdallah medical city

¹²Dental assistant, Primary health care center in Alzimah

¹³Muhayil Specialized Dental Center, Dental technician

¹⁴Dental technician at Altaneem PHC

¹⁵TECHNICIAN RADIOLOGY TECHNOLOGY, KING FAHAD GENERAL HOSPITAL JEDDAH

ABSTRACT

Odontogenic cysts represent a significant portion of jaw pathologies encountered in clinical practice. These cystic lesions originate from epithelial remnants associated with the development of teeth and are typically classified into inflammatory or developmental types. Common examples include radicular cysts, dentigerous cysts, and odontogenic keratocysts (OKCs), the latter being known for its aggressive behavior and high recurrence rate. The prevalence and diverse presentations of odontogenic cysts highlight the importance of accurate diagnosis and effective management [1].

The clinical presentation of odontogenic cysts varies widely, from asymptomatic radiographic findings to significant jaw swelling, pain, or even pathological fractures. Diagnosis is often challenging due to the overlap in radiographic appearances among different cyst types, as well as with other odontogenic and non-odontogenic lesions. This underscores the pivotal role of imaging modalities in identifying and characterizing these lesions. Beyond diagnosis, imaging is integral to treatment planning, monitoring disease progression, and evaluating post-treatment outcomes.

Historically, conventional imaging modalities such as intraoral radiography and panoramic radiography have been the mainstay for the evaluation of odontogenic cysts. These tools offer valuable initial insights but are often limited by their two-dimensional nature and inability to capture the complete extent of a lesion. The advent of advanced imaging techniques, including cone-beam computed tomography (CBCT), computed tomography (CT), and magnetic resonance imaging (MRI), has revolutionized the diagnostic landscape. These modalities provide three-dimensional views, detailed anatomical relationships, and improved characterization of lesion properties [2,3].

Advancements in imaging technology, coupled with the integration of artificial intelligence (AI) and low-dose imaging protocols, are further enhancing diagnostic precision while minimizing patient risks. These innovations promise to address key challenges, such as differentiating between cystic and solid lesions, assessing proximity to critical structures, and planning minimally invasive interventions [4].

This review aims to explore the role of imaging modalities in the diagnosis and management of odontogenic cysts. By synthesizing current literature, we examine the strengths and limitations of each imaging technique, discuss their clinical applications, and consider emerging technologies that are shaping the future of maxillofacial diagnostics and treatment.

Review:

Conventional Imaging Modalities

1. Intraoral Radiography

Intraoral radiography, including periapical and occlusal views, remains a mainstay in routine dental practice due to its accessibility and high resolution for localized areas. These images are particularly useful in detecting small periapical lesions, often presenting as well-defined radiolucencies [3]. For instance, a periapical radiograph can effectively detect a radicular cyst associated with a non-vital tooth.

However, the limitations of intraoral radiography lie in its two-dimensional nature, which can obscure the true extent and depth of lesions. Additionally, overlapping anatomical structures, such as the zygomatic arch or maxillary sinus, may hinder accurate interpretation [4].

2. Panoramic Radiography

Panoramic radiographs offer a broader view of the maxillofacial region, making them ideal for evaluating larger lesions and their relationship to adjacent structures. They are particularly useful in identifying dentigerous cysts, which often appear as unilocular radiolucencies surrounding the crown of an unerupted tooth [5].

Despite their utility, panoramic radiographs have limitations, including geometric distortion and reduced resolution compared to intraoral radiographs. As such, they are typically used as a screening tool rather than for detailed lesion characterization.

Advanced Imaging Modalities

1. Cone-Beam Computed Tomography (CBCT)

Mohammed Ahmed Alhudiry, Mohammed Ibrahim Fageeh, Abdullah Hussain Alasmari, Mohammed Ali Ahmed Assiri, Faisal Hadi Abdullah Asiri, Abdullah Mohammed Alsayed, Mohammed Ibrahim Muidh Althui, Nahla Atia Nasser Albarakati, Hasnaa Eshaq Haroun Hawsawi, Hashem Alawi Melebary, Kholood Harisi, Ohood Abdaraboh Alhabashi, Nasir Abduh Mana Al Ahmari, Amal Saleh Alhiani, Wafaa Mohammed Alomari

CBCT has revolutionized the field of maxillofacial imaging by providing three-dimensional, high-resolution images at a relatively lower radiation dose than conventional computed tomography (CT). It allows for precise measurement of lesion dimensions, internal content, and proximity to vital structures such as the inferior alveolar nerve and maxillary sinus [6].

For odontogenic keratocysts, CBCT plays a crucial role in evaluating cortical bone expansion and perforation, both of which are significant in determining the aggressiveness of the lesion. Studies have demonstrated that CBCT can identify features such as scalloped margins and multilocular appearances, which are characteristic of OKCs [7].

2. Computed Tomography (CT)

CT imaging is particularly valuable in cases where soft tissue involvement or complex bone destruction is suspected. Although CBCT has largely replaced CT for routine dental imaging, CT remains superior in visualizing soft-tissue contrast. For instance, in cases of infected cysts or suspected malignancy, contrast-enhanced CT can delineate the extent of soft tissue invasion [8].

However, the higher radiation dose associated with CT compared to CBCT limits its use, especially in younger patients and those requiring serial imaging.

3. Magnetic Resonance Imaging (MRI)

MRI provides exceptional soft-tissue contrast, making it a powerful tool in differentiating between cystic and solid lesions. Unlike CT and CBCT, MRI does not involve ionizing radiation, making it suitable for pediatric and pregnant patients. For odontogenic cysts, T1- and T2-weighted sequences can help identify fluid content, septations, and potential adjacent soft-tissue involvement [9].

MRI is particularly useful in distinguishing odontogenic cysts from tumors, such as ameloblastomas or odontogenic carcinomas, which may exhibit overlapping features on CT or CBCT [10].

Comparative Analysis of Imaging Modalities

A comparative analysis of the imaging modalities highlights their strengths and weaknesses in the context of odontogenic cyst diagnosis and management.

Imaging Modality	Strengths	Limitations
Intraoral Radiography	High resolution, low cost	Limited field of view, no 3D information
Panoramic Radiography	Wide field of view, accessible	Distortion, reduced resolution
CBCT	High spatial resolution, 3D imaging	Radiation exposure, higher cost

Imaging Modality	Strengths	Limitations
CT	Superior soft-tissue and bone contrast	High radiation dose, limited availability
MRI	Excellent soft-tissue contrast, no radiation	High cost, limited availability

Applications in Management

1. Diagnosis

Imaging aids in the initial differentiation of odontogenic cysts based on their radiographic features. For example:

Radicular cysts typically present as well-defined, unilocular radiolucencies at the apex of a non-vital tooth [11].

Dentigerous cysts are often associated with impacted teeth, exhibiting unilocular radiolucency surrounding the crown [12].

OKCs display multilocular, scalloped radiolucencies with a tendency for cortical perforation [13].

2. Treatment Planning

Imaging facilitates treatment planning by delineating lesion boundaries, internal structures, and proximity to vital anatomical features. For instance, CBCT can guide enucleation of a dentigerous cyst while avoiding damage to the inferior alveolar nerve [14].

3. Post-Treatment Follow-Up

Postoperative imaging is essential for assessing treatment outcomes and detecting recurrence. For OKCs, CBCT is preferred due to its sensitivity in identifying small residual lesions that may not be visible on conventional radiographs [15].

Emerging Technologies

Artificial Intelligence (AI)

AI-powered diagnostic tools are increasingly being integrated into imaging workflows. Machine learning algorithms can enhance lesion detection and classification, reducing diagnostic variability among clinicians. For example, AI models trained on CBCT datasets have demonstrated high accuracy in identifying odontogenic cysts and differentiating them from tumors [16].

Low-Dose Imaging Protocols

To address concerns about radiation exposure, particularly in pediatric populations, low-dose CBCT protocols have been developed. These protocols maintain diagnostic image quality while significantly reducing patient radiation exposure [16].

3D Printing and Guided Surgery

The integration of CBCT data with 3D printing technology has enabled the creation of surgical guides for precise lesion enucleation. This approach minimizes intraoperative risks and improves surgical outcomes [16].

Discussion and Future Directions

While imaging modalities have significantly improved the diagnosis and management of odontogenic cysts, challenges remain. The interpretation of imaging findings requires a high level of expertise, and misdiagnosis can lead to inappropriate treatment. Additionally, the cost and availability of advanced imaging technologies may limit their use in resource-constrained settings.

Future research should focus on the development of standardized imaging protocols and the integration of multimodal imaging techniques to enhance diagnostic accuracy. The role of AI in predictive modeling and treatment planning also warrants further exploration.

Conclusion

Imaging modalities are indispensable in the accurate diagnosis and effective management of odontogenic cysts. These cystic lesions, while often benign, can lead to significant complications if not promptly identified and treated. Conventional imaging techniques such as intraoral radiography and panoramic radiography remain valuable tools for initial screening and diagnosis, particularly for smaller or more localized lesions. However, their limitations in providing comprehensive three-dimensional information have been addressed by advanced imaging modalities like CBCT, CT, and MRI.

Among these, CBCT has emerged as the gold standard for the evaluation of odontogenic cysts. Its ability to provide high-resolution, three-dimensional images makes it ideal for assessing lesion size, extent, and relationship to vital anatomical structures. CT and MRI also play critical roles, particularly in complex cases where soft-tissue contrast or detailed anatomical visualization is required. While CT is preferred for its superior bone and soft tissue imaging, MRI offers an alternative without ionizing radiation, making it especially valuable in pediatric and pregnant patients.

The integration of advanced technologies such as AI and 3D printing is transforming the field, enabling more precise lesion detection, classification, and treatment planning. Low-dose imaging protocols further enhance patient safety, addressing concerns about cumulative radiation exposure, especially in individuals requiring long-term monitoring.

Despite these advancements, challenges remain. Variability in diagnostic interpretation, cost barriers, and limited access to advanced imaging technologies in resource-constrained settings highlight areas for further development. Collaborative

efforts to standardize imaging protocols and integrate multimodal imaging approaches can enhance diagnostic accuracy and streamline clinical workflows.

References

- Ackermann G, Cohen MA, Altini M. The paradental cyst: a clinicopathologic study of 50 cases. *Oral Surg Oral Med Oral Pathol.* 1987 Sep;64(3):308-12.
- Bilodeau EA, Collins BM. Odontogenic Cysts and Neoplasms. *Surg Pathol Clin.* 2017 Mar;10(1):177-222.
- Carrillo C, Peñarocha M, Bagán JV, Vera F. Relationship between histological diagnosis and evolution of 70 periapical lesions at 12 months, treated by periapical surgery. *J Oral Maxillofac Surg.* 2008 Aug;66(8):1606-9.
- Chirapathomsakul D, Sastravaha P, Jansitsyanont P. A review of odontogenic keratocysts and the behavior of recurrences. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2006 Jan;101(1):5-9; discussion 10.
- Chrcanovic BR, Reis BM, Freire-Maia B. Paradental (mandibular inflammatory buccal) cyst. *Head Neck Pathol.* 2011 Jun;5(2):159-64.
- Cohen DA, Neville BW, Damm DD, White DK. The lateral periodontal cyst. A report of 37 cases. *J Periodontol.* 1984 Apr;55(4):230-4.
- Daley TD, Wysocki GP. The small dentigerous cyst. A diagnostic dilemma. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 1995 Jan;79(1):77-81.
- de Souza LB, Gordón-Núñez MA, Nonaka CF, de Medeiros MC, Torres TF, Emiliano GB. Odontogenic cysts: demographic profile in a Brazilian population over a 38-year period. *Med Oral Patol Oral Cir Bucal.* 2010 Jul 01;15(4):e583-90.
- Dhawan P, Kochhar GK, Chachra S, Advani S. Eruption cysts: A series of two cases. *Dent Res J (Isfahan).* 2012 Sep;9(5):647-50.
- Fowler CB, Brannon RB, Kessler HP, Castle JT, Kahn MA. Glandular odontogenic cyst: analysis of 46 cases with special emphasis on microscopic criteria for diagnosis. *Head Neck Pathol.* 2011 Dec;5(4):364-75.
- Johnson NR, Gannon OM, Savage NW, Batstone MD. Frequency of odontogenic cysts and tumors: a systematic review. *J Investig Clin Dent.* 2014 Feb;5(1):9-14.
- Molven O, Halse A, Grung B. Incomplete healing (scar tissue) after periapical surgery--radiographic findings 8 to 12 years after treatment. *J Endod.* 1996 May;22(5):264-8.
- Rioux-Forker D, Deziel AC, Williams LS, Muzaffar AR. Odontogenic Cysts and Tumors. *Ann Plast Surg.* 2019 Apr;82(4):469-477.
- Thosaporn W, Iamaroon A, Pongsiriwet S, Ng KH. A comparative study of epithelial cell proliferation between the odontogenic keratocyst, orthokeratinized odontogenic cyst, dentigerous cyst, and ameloblastoma. *Oral Dis.* 2004 Jan;10(1):22-6.
- Titinchi F, Nortje CJ. Keratocystic odontogenic tumor: a recurrence analysis of clinical and radiographic parameters. *Oral Surg Oral Med Oral Pathol Oral Radiol.* 2012 Jul;114(1):136-42.
- Weber AL. Imaging of cysts and odontogenic tumors of the jaw. Definition and classification. *Radiol Clin North Am.* 1993 Jan;31(1):101-20.