

Effects of Nurse-Led Selfcare Interventions on Health Outcomes among People with Heart Failure

Majed Haumed Al Zaidy¹, Sarah Talaq Almotairi², Afaf Abdullah Al-Dawsari³, Maha Othman Badri⁴, Lamia Nasser Alruwaili⁵, Latifa Rashid Almuteri⁶, Faizah Ateeg ALBoeinain⁷, Mona Hamdan Nahar Albelaasi⁵, Bashayer Ayed Farhan AlAnzi⁸, Alia Awad Farhan Al-Anzi⁸, Abdulrahman Bahri Madhan Al-Omrani⁹, Ehsan sayeg alanizi¹⁰

1. Nursing, Riyadh Specialised Dental Center, Riyadh, KSA
2. Nursing Technician, Laban Center 2, Riyadh, KSA
3. Nursing Technician, King Fahad Medical City, Riyadh, KSA
4. Nursing Specialist, Seha Virtual Hospital, Riyadh, KSA
5. Nursing Technician, Qurayyat General Hospital, Qurayyat, KSA
6. Nursing, King Fahad Health Center, Riyadh, KSA
7. Nursing, Markaz zahrat albdye, Riyadh, KSA
8. Nursing Technician, Specialized Dental Hospital, Arar, KSA
9. Nursing, Tabuk Health Cluster, King Fahad Specialist Hospital in Tabuk, KSA
10. Nursing technician, Al-Qurayyat General Hospital, Al Qurayyat, KSA

ABSTRACT

This study investigates the impact of nurse-led self-care interventions on health outcomes in patients with heart failure (HF), a condition characterized by the heart's inability to pump blood effectively, resulting in significant morbidity, mortality, and healthcare burdens. Given the complexities of HF management, including medication adherence and lifestyle modifications, effective self-care is crucial. However, many patients face challenges in self-care due to factors such as lack of knowledge, emotional distress, and insufficient support. This research highlights the pivotal role of nurses in providing education and support to empower patients, enhance health literacy, and foster engagement in self-management practices. Through comprehensive assessments, nurses can tailor interventions to meet individual needs, addressing barriers such as social determinants of health and mental health conditions. The study employs a robust evaluation framework encompassing clinical

Effects of Nurse-Led Selfcare Interventions on Health Outcomes among People with Heart Failure metrics, such as hospital readmission rates and changes in ejection fraction, alongside quality of life assessments using validated instruments like the Kansas City Cardiomyopathy Questionnaire. By analyzing these outcomes pre- and postintervention, the study aims to elucidate the effectiveness of nurse-led self-care strategies in improving health outcomes for HF patients. Furthermore, it emphasizes the importance of patient feedback and engagement in the design and implementation of interventions, recognizing that understanding patient barriers and incorporating digital health tools can enhance adherence to self-care practices. The findings are expected to contribute to the growing body of evidence supporting nurse-led interventions as a viable strategy to improve health outcomes in heart failure management, ultimately leading to reduced hospitalizations, enhanced quality of life, and better overall patient empowerment in managing their health. This research underscores the necessity for innovative, nurse-driven approaches to address the multifaceted challenges faced by heart failure patients, aiming to bridge the gap between clinical guidelines and personalized care.

KEYWORDS: heart failure, self-care, nursing.

1. Introduction

Heart failure (HF) represents a significant public health challenge, affecting millions of individuals worldwide. It occurs when the heart is unable to pump sufficiently to maintain blood flow to meet the body's needs, resulting in a complex clinical syndrome characterized by symptoms such as fatigue, shortness of breath, and fluid retention. [1] This condition can arise from various underlying causes, including coronary artery disease, hypertension, and valvular heart disease, and often leads to frequent hospitalizations, reduced quality of life, and increased mortality. Given the progressive nature of HF, effective self-care management becomes critical for patients, enabling them to manage their symptoms and maintain their health. [2]

Self-care in heart failure encompasses a range of behaviors and practices that individuals can engage in to take charge of their health. This includes daily monitoring of weight, adherence to medication regimens, managing dietary restrictions, and recognizing early signs of deterioration. [3] Nevertheless, many patients with HF struggle with self-care due to a myriad of factors, including a lack of knowledge, emotional distress, and insufficient support from healthcare providers. As the burden of heart failure continues to grow, innovative strategies to enhance self-care and improve health outcomes are paramount. One such strategy involves nurse-led self-care interventions, which have been shown to empower patients, enhance their understanding of the condition, and improve adherence to treatment plans. [4]

The role of nurses in heart failure management is increasingly recognized as vital for fostering self-care behaviors. Nurse-led interventions focus on education, motivation,

Majed Haumed Al Zaidy¹, Sarah Talaq Almotairi², Afaf Abdullah Al-Dawsari³, Maha Othman Badri⁴, Lamia Nasser Alruwaili⁵, Latifa Rashid Almuteri⁶, Faizah Ateeg ALBoeinain⁷, Mona Hamdan Nahar Albelaasi⁵, Bashayer Ayed Farhan AlAnzi⁸, Alia Awad Farhan Al-Anzi⁸, Abdulrahman Bahri Madhan Al-Omran⁹, Ehsan sayeg alanizi¹⁰

and support, addressing the unique needs of each patient. By providing individualized care plans, nursing staff can help patients understand the importance of weighing themselves daily, recognizing changes in symptoms, and coordinating medication schedules. [5] Moreover, these interventions often incorporate motivational interviewing techniques, which encourage patients to express their concerns and preferences, ultimately leading to a more engaged and proactive approach to health management.

The effectiveness of nurse-led self-care interventions is not merely limited to physiological improvements; they also contribute to the psychological well-being of patients. Heart failure is often accompanied by anxiety and depression, which can significantly impact a patient's ability to manage their condition effectively. By fostering a supportive therapeutic relationship, nurses can help mitigate these feelings, promoting a sense of control and self-worth among patients as they navigate their healthcare journey.[6] Regular follow-ups and check-ins allow nurses to provide encouragement, track progress, and adjust care plans as needed, reinforcing positive behaviors and outcomes.

Moreover, the integration of technology into nurse-led self-care interventions offers new avenues for enhancing patient engagement. Tools such as mobile health applications, telehealth consultations, and remote monitoring devices provide patients with real-time feedback and support, helping them stay connected to their healthcare team. This technological integration further empowers individuals to take an active role in their self-care, promoting autonomy and reinforcing the importance of adherence to treatment regimens. [7] Objectives:

The main objectives of this review are to:

1. Evaluate how nurse-led self-care interventions improve the self-care abilities of individuals with heart failure.
2. Assess the impact of interventions on patient care.
3. Analyze the cost-effectiveness of nurse-led self-care interventions in managing heart failure compared to standard care practices in terms of healthcare utilization costs.

The Role of Nurses in Patient Care:

One of the primary roles of nurses in heart failure care is patient education. Nurses equip patients and their families with crucial information regarding heart failure management, including lifestyle modifications, symptom recognition, and the importance of medication adherence. Educational interventions led by nurses have been shown to result in a deeper understanding of the condition and its management among patients. Stakeholders within a farmed study titled "Effects of Nurse-Led Self-

Effects of Nurse-Led Selfcare Interventions on Health Outcomes among People with Heart Failure
Care Interventions on Health Outcomes Among People with Heart Failure" highlighted the positive impacts of structured educational programs instituted by nursing professionals. Nurses employ various strategies to enhance health literacy, using teach-back methods, visual aids, and printed materials, thus empowering patients to take charge of their health. [8]

Furthermore, nurses are integral in conducting comprehensive assessments that inform personalized care plans. This involves not only evaluating the physical symptoms of heart failure but also understanding the psychosocial aspects that impact a patient's ability to manage their condition. Through thorough assessments, nurses can identify barriers to self-care, such as social determinants of health, mental health conditions, and cognitive limitations. By addressing these factors, nurses can customize interventions that cater to the unique needs of individuals with heart failure. This holistic approach fosters a supportive environment that encourages patient engagement and self-efficacy. [9]

The implementation of nurse-led self-care interventions is another critical aspect of nursing in heart failure care. Such interventions may include structured monitoring of symptoms, encouraging regular physical activity, and facilitating adherence to diet and medication regimens. Nurses often utilize telehealth technology to maintain regular communication with patients, allowing for real-time assessments and timely intervention in case of symptom exacerbation. For instance, studies have demonstrated that telemonitoring systems led by nurses can reduce hospitalization rates and improve quality of life for heart failure patients. In this regard, the nurse's role transcends traditional care limits, employing innovative strategies to foster improved health outcomes. [10]

Additionally, the emotional support provided by nurses is vital in managing heart failure. The chronic nature of HF can lead to feelings of anxiety, depression, and social isolation in patients. Nurses are often the primary caregivers who offer empathetic listening and counseling, helping patients navigate their feelings about living with a chronic illness. By establishing a trusting nurse-patient relationship, nurses can promote mental well-being and encourage positive behavioral changes that contribute to better health outcomes. This psycho-social support complements the clinical care and serves to enhance the overall quality of life for individuals facing the challenges of heart failure. [11]

Implementation of Nurse-Led Self-Care Interventions

A growing body of evidence advocates for the implementation of nurse-led self-care interventions, which empower patients to actively participate in managing their condition. This approach places nurses in a pivotal role, bridging the gap between clinical guidelines and individualized patient care. The aim of this study is to evaluate the effects of these interventions on health outcomes among patients with heart failure. [12]

Majed Haumed Al Zaidy¹, Sarah Talaq Almotairi², Afaf Abdullah Al-Dawsari³, Maha Othman Badri⁴, Lamia Nasser Alruwaili⁵, Latifa Rashid Almuteri⁶, Faizah Ateeg ALBoeinain⁷, Mona Hamdan Nahar Albelaasi⁵, Bashayer Ayed Farhan AlAnzi⁸, Alia Awad Farhan Al-Anzi⁸, Abdulrahman Bahri Madhan Al-Omrani⁹, Ehsan sayeg alanizii¹⁰

The first step in the implementation of nurse-led self-care interventions is conducting a comprehensive assessment of each patient's individual needs, preferences, and existing knowledge regarding heart failure management. This assessment allows nurses to identify specific areas where education and support are required. [13] For instance, patients may need assistance in recognizing early signs of decompensation or understanding the critical importance of adhering to dietary restrictions and medication regimens. Additionally, factors such as social support, mental health status, and health literacy significantly affect individuals' ability to conduct effective self-care. By understanding these dimensions, nurses can tailor self-care plans that resonate with each patient's unique circumstances and encourage adherence to necessary lifestyle modifications. [14]

Education stands at the forefront of nurse-led self-care interventions. Nurses are tasked with not only disseminating essential information about heart failure but also ensuring that patients comprehend this information. Interactive educational sessions that engage patients through discussions, role-playing, and problem-solving can significantly enhance understanding and retention. Providing resources like brochures, videos, and digital apps can reinforce the learning process and offer ongoing references for patients. Additionally, teaching strategies that include motivational interviewing can empower patients by fostering a sense of autonomy in their self-care practices, enhancing their confidence and ability to make informed decisions regarding their health. [15]

Moreover, establishing individualized self-care goals is crucial in this intervention process. Using a goal-setting approach aligns with the principles of patient-centered care, promoting engagement and accountability in managing their health. For instance, goals can range from daily weight monitoring to achieving a certain level of physical activity based on individual capacity and health status. Regular follow-up appointments play an essential role in facilitating progress towards these goals. During these visits, nurses can review patients' self-care practices, address challenges encountered, and adjust the care plan as necessary. This ongoing support not only reinforces learning but also helps patients feel less isolated as they navigate their health journeys. [16]

In many cases, nurse-led interventions also incorporate collaborative care models, where multidisciplinary team members, including dietitians, pharmacists, and social workers, contribute to comprehensive care. A coordinated approach ensures that all aspects of the patient's health and well-being are considered, permitting the identification of potential barriers to adherence to self-care strategies. [17] Involving family members or caregivers in the educational process can further augment these efforts by creating a support network that reinforces self-care practices at home.

The evaluation of nurse-led self-care interventions should include both qualitative and quantitative measures to assess their impact comprehensively. Quantitative measures may include clinical outcomes such as hospitalization rates, medication adherence rates, and the frequency of emergency department visits. [18] Qualitative feedback from patients regarding their experiences and perceptions of the interventions can illuminate pathways for enhancing care quality and meeting patients' changing needs. Engagement in follow-up surveys or interviews can deepen the understanding of how self-care practices affect patients' daily lives and satisfaction with their health outcomes. [19]

Ultimately, the success of nurse-led self-care interventions in heart failure management hinges upon the incorporation of evidence-based practice, ongoing education, and a patient-centered approach to care. As healthcare delivery increasingly shifts toward models that promote self-management, the essential role of nurses as educators, motivators, and advocates will become increasingly significant. Enhancing patients' self-efficacy through structured support systems and collaborative care will not only lead to improved health outcomes but also address the emotional and psychological aspects of living with a chronic condition like heart failure. Future research should continue to explore various strategies to implement and sustain nurse-led self-care interventions while determining best practices for patient education and engagement. Through these endeavors, we can pave the way toward optimizing care and nurturing resilience in those challenged by heart failure. [20]

Impact of Nurse-Led Self-Care Interventions:

The assessment of health outcomes in patients with heart failure (HF) following nurse-led self-care interventions is multifaceted and necessitates a robust framework for evaluation metrics. Heart failure, characterized by the heart's inability to pump adequately and meet the body's demands, often leads to significant morbidity and mortality. As a chronic condition, it imposes substantial burdens on the healthcare system, making effective management critical. Nurse-led self-care interventions aim to empower patients by teaching them skills to manage their condition, which is vital in improving health outcomes. To accurately evaluate the effectiveness of these interventions, specific metrics need to be established, taking into account clinical, behavioral, and patient-reported outcomes. [21] One of the primary clinical metrics is the rate of hospital readmissions. In heart failure patients, frequent hospitalizations are indicative of worsening health and can be linked to poor self-management, medication non-adherence, or lack of knowledge about lifestyle changes. Studies have shown that effective self-care interventions, particularly those led by nurses, can reduce readmission rates significantly. Tracking the frequency of these readmissions pre- and post-intervention provides a clear picture of the intervention's impact on overall health outcomes. [22]

Majed Haumed Al Zaidy¹, Sarah Talaq Almotairi², Afaf Abdullah Al-Dawsari³, Maha Othman Badri⁴, Lamia Nasser Alruwaili⁵, Latifa Rashid Almuteri⁶, Faizah Ateeg ALBoeinain⁷, Mona Hamdan Nahar Albelaasi⁵, Bashayer Ayed Farhan AlAnzi⁸, Alia Awad Farhan Al-Anzi⁸, Abdulrahman Bahri Madhan Al-Omrani⁹, Ehsan sayeg alanizii¹⁰

Another important clinical metric is the measurement of morbidity, which can be assessed by evaluating changes in ejection fraction (EF) or assessing the New York Heart Association (NYHA) functional classification. Both metrics provide crucial insights into the cardiac function and functional capacity of patients. For instance, improvements in EF or a shift to a lower NYHA class (indicating less severe heart failure) after implementing nurse-led interventions can substantiate the program's effectiveness in enhancing patients' health. [23]

Additionally, quality of life (QoL) metrics must be integrated into the evaluation framework. Tools such as the Kansas City Cardiomyopathy Questionnaire (KCCQ) or the Minnesota Living with Heart Failure Questionnaire (MLHFQ) are standardized and validated instruments that help gauge a patient's QoL related to heart failure specifically. By analyzing scores from these instruments before and after nurse-led interventions, researchers can assess the subjective improvements in patients' daily living and emotional well-being that may result from effective selfmanagement strategies imparted by nursing staff. [24]

Behavioral changes are also crucial metrics in evaluating nurse-led self-care interventions. Self-efficacy, which refers to a patient's belief in their ability to manage their health, can be assessed using the Self-Efficacy for Managing Chronic Disease 6-Item Scale (SEMCD). Additionally, adherence rates to prescribed medications, dietary recommendations, and exercise regimens should be closely monitored. Self-reported adherence, coupled with pharmacy refill data where applicable, offers an invaluable perspective on whether patients are not only learning but also implementing self-care practices in their daily lives. [25]

Moreover, patient satisfaction should be evaluated as an integral component of the overall health outcome framework. The experiences and perceptions of patients regarding the care they receive can significantly influence their engagement in selfcare practices. Surveys that measure satisfaction with care, clarity of health education received, and perceived support from nursing staff can help determine the subjective effectiveness of the interventions. High levels of satisfaction often correlate with increased adherence and, consequently, improved health outcomes. [26]

Finally, cost-effectiveness analysis should also be considered as part of the evaluation metrics. The economic impact of nurse-led self-care interventions can be evaluated by measuring healthcare utilization costs, including hospitalizations and emergency visits, against the outcomes achieved. This analysis will illustrate the financial benefits of these interventions, making the case for their broader implementation within the healthcare system. [27]

Patient Feedback and Engagement with Self-Care Interventions

Patient feedback and engagement are crucial components in the effective implementation of self-care interventions, particularly for individuals living with chronic conditions such as heart failure. Self-care, the ability of individuals to manage their health through a combination of knowledge, skills, and behaviors, becomes particularly important in settings where patients are faced with the complexities of managing symptoms and adhering to treatment regimens. Within the context of heart failure, where adherence to medication, lifestyle modifications, and symptom recognition are pivotal, nurse-led self-care interventions have emerged as a vital strategy to promote better health outcomes. [28]

For interventions to be successful, they must be designed with the patient's perspective at the forefront. This involves a thorough understanding of the barriers patients face in implementing self-care strategies. Patients may encounter various obstacles, including financial constraints, lack of social support, and psychological challenges like anxiety or depression. Addressing these barriers within self-care interventions is essential for promoting greater engagement. For example, offering financial assistance programs or access to community resources can empower patients to manage their condition more effectively. Furthermore, fostering a supportive environment where emotions and concerns can be openly discussed may alleviate some psychological burdens, enhancing patients' ability to adhere to selfcare practices. [29]

The integration of digital health tools, such as mobile applications and telehealth platforms, has opened new avenues for patient engagement in self-care. These tools facilitate real-time communication between healthcare providers and patients, enabling the timely exchange of information. Such interactions can lead to more immediate adjustments in care plans and more responsive support for patients facing difficulties. Through gamification and interactive features, mobile applications can also encourage patients to track their progress, set health goals, and receive personalized feedback, which can further motivate them to take ownership of their health. [30]

However, while these technologies offer great potential, it is crucial that their implementation is guided by patient preferences and accessibility considerations. Not all patients may feel comfortable using digital tools, and disparities in technology access can pose significant challenges. Nurses must remain attentive to these disparities and strive to offer alternate methods of engagement to ensure that all patients have the opportunity to benefit from self-care interventions. [31]

The role of nurses in facilitating patient feedback cannot be overstated. Nurses serve as the primary point of contact within the healthcare system, making them ideally positioned to advocate for patients and to cultivate an environment conducive to open dialogue. Through regular check-ins, educational sessions, and follow-up visits, nurses can solicit patient input regarding their experiences with self-care interventions. This dialogue should not be a one-time event but rather an ongoing process that continuously informs and refines care strategies. [32]

Majed Haumed Al Zaidy1, Sarah Talaq Almotairi2, Afaf Abdullah Al-Dawsari3, Maha Othman Badri4, Lamia Nasser Alruwaili5, Latifa Rashid Almuteri6, Faizah Ateeg ALBoeinain7, Mona Hamdan Nahar Albelaasi5, Bashayer Ayed Farhan AlAnzi8, Alia Awad Farhan Al-Anzi8, Abdulrahman Bahri Madhan Al-Omrani9, Ehsan sayeg alanizi10

Understanding that the effectiveness of nurse-led self-care interventions is rooted in patient engagement and feedback allows for a more holistic approach to heart failure management. By fostering a culture of collaboration between patients and healthcare providers, and by consistently tailoring interventions based on patient responses, the healthcare system can ultimately improve health outcomes and enhance the quality of life for individuals living with heart failure. As research in this field continues to evolve, it is imperative that strategies remain adaptable and that patient voices remain central in the development and implementation of self-care interventions. This alignment not only leads to better health outcomes but also builds a resilient and empowered patient population capable of effectively managing their chronic conditions. [33]

2. Conclusion:

In conclusion, this study underscores the critical role of nurse-led self-care interventions in managing heart failure, a condition that poses significant challenges to patients and healthcare systems alike. By equipping patients with essential knowledge and skills, nurses can empower individuals to take an active role in their health management, leading to improved health outcomes and enhanced quality of life. The evidence highlights that tailored educational programs, comprehensive assessments, and the integration of digital health tools significantly contribute to patient engagement and adherence to self-care practices. Moreover, addressing barriers such as social support and mental health is crucial for fostering effective self-management. As healthcare continues to evolve, the implementation of innovative, nurse-led strategies will be vital in bridging the gap between clinical guidelines and personalized care, ultimately reducing hospitalizations and improving overall patient well-being in heart failure management. Future research should focus on refining these interventions and exploring their long-term impacts on patient health outcomes.

References

- Von Lueder T.G., Agewall S. The burden of heart failure in the general population: A clearer and more concerning picture. *J. Thorac. Dis.* 2018;10:S1934–S1937. doi: 10.21037/jtd.2018.04.153. [DOI] [PMC free article] [PubMed] [Google Scholar]
- Savarese G., Lund L.H. Global public health burden of heart failure. *Card. Fail. Rev.* 2017;3:7–11. doi: 10.15420/cfr.2016:25:2. [DOI] [PMC free article] [PubMed] [Google Scholar]
- Go A.S., Mozaffarian D., Roger V.L., Benjamin E.J., Berry J.D., Blaha M.J., Dai S., Ford E.S., Fox C.S., Franco S., et al. Heart disease and stroke statistics—2014 update: A report from the American Heart Association. *Circulation.* 2014;129:e28–e292. doi: 10.1161/01.cir.0000441139.02102.80. [DOI] [PMC free article] [PubMed] [Google Scholar]
- Lesyuk W., Kriza C., Kolominsky-Rabas P. Cost-of-illness studies in heart failure: A systematic review 2004-2016. *BMC Cardiovasc. Disord.* 2018;18 doi: 10.1186/s12872-018-0815-3. [DOI] [PMC free article] [PubMed] [Google Scholar]

- Effects of Nurse-Led Selfcare Interventions on Health Outcomes among People with Heart Failure
 Salahodinkolah M.K., Ganji J., Moghadam S.H., Shafipour V., Jafari H., Salari S. Educational intervention for improving self-care behaviors in patients with heart failure: A narrative review. *J. Nurs. Midwifery Sci.* 2020;7:60–68. doi: 10.4103/JNMS.JNMS_19_19. [DOI] [Google Scholar]
- Ba H.M., Son Y., Lee K., Kim B.-H. Transitional Care Interventions for Patients with Heart Failure: An Integrative Review. *Int. J. Environ. Res. Public Health.* 2020;17:2925. doi: 10.3390/ijerph17082925. [DOI] [PMC free article] [PubMed] [Google Scholar]
- Greene S.J., Fonarow G.C., Vaduganathan M., Khan S.S., Butler J., Gheorghiade M. The vulnerable phase after hospitalization for heart failure. *Nat. Rev. Cardiol.* 2015;12:220–229. doi: 10.1038/nrcardio.2015.14. [DOI] [PubMed] [Google Scholar]
- Lee C.S., Mudd J.O., Hiatt S.O., Gelow J.M., Chien C., Riegel B. Trajectories of heart failure self-care management and changes in quality of life. *Eur. J. Cardiovasc. Nurs.* 2015;14:486–494. doi: 10.1177/1474515114541730. [DOI] [PubMed] [Google Scholar]
- McMurray J.J., Adamopoulos S., Anker S.D., Auricchio A., Böhm M., Dickstein K., Falk V., Filippatos G., Fonseca C., Gomez-Sanchez M.A., et al. ESC Committee for Practice Guidelines. ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure 2012: The task force for the diagnosis and treatment of acute and chronic heart failure 2012 of the European Society of Cardiology. Developed in collaboration with the Heart Failure Association (HFA) of the ESC. *Eur. Heart J.* 2012;33:1787–1847. doi: 10.1093/eurjhf/hfs105. [DOI] [PubMed] [Google Scholar]
- Toukhsati S.R., Driscoll A., Hare D.L. Patient self-management in chronic heart failure—Establishing concordance between guidelines and practice. *Card. Fail. Rev.* 2015;1:128–131. doi: 10.15420/cfr.2015.1.2.128. [DOI] [PMC free article] [PubMed] [Google Scholar]
- Jonkman N.H., Westland H., Groenwold R.H.H., Ågren S., Atienza F., Blue L., de la Bruggink-André Porte P.W.F., Darren A DeWalt D.A., Hebert P.L., Heisler M., et al. Do self-management interventions work in patients with heart failure? An individual patient data meta-analysis. *Circulation.* 2016;133:1189–1198. doi: 10.1161/CIRCULATIONAHA.115.018006. [DOI] [PMC free article] [PubMed] [Google Scholar]
- Herber O.R., Atkins L., Störk S., Wilm S. Enhancing self-care adherence in patients with heart failure: A study protocol for developing a theory-based behaviour change intervention using the COM-B behaviour model (ACHIEVE study) *BMJ Open.* 2018;8:e025907. doi: 10.1136/bmjopen-2018025907. [DOI] [PMC free article] [PubMed] [Google Scholar]
- Seid M.A., Abdela O.A., Zeleke E.G. Adherence to self-care recommendations and associated factors among adult heart failure patients. From the patients' point of view. *PLoS ONE.* 2019;14:e0211768. doi: 10.1371/journal.pone.0211768. [DOI] [PMC free article] [PubMed] [Google Scholar]
- Jaarsma T., Cameron J., Riege B., Stromberg A. Factors related to self-care in heart failure patients according to the middle-range theory of self-care of chronic illness: A literature update. *Curr. Heart Fail. Rep.* 2017;14:71–77. doi: 10.1007/s11897-017-0324-1. [DOI] [PMC free article] [PubMed] [Google Scholar]
- Francis L.P., Spaulding E., Turkson-Ocran R.A., Allen J. Randomized trials of nurse-delivered interventions in weight management research: A systematic review. *West. J. Nurs. Res.* 2017;39:1120–1150. doi: 10.1177/0193945916686962. [DOI] [PMC free article] [PubMed] [Google Scholar]
- Rasmusson K., Flattery M., Baas L.S. American association of heart failure nurses position paper on educating patients with heart failure. *Heart Lung.* 2015;44:173–177. doi: 10.1016/j.hrtlng.2015.01.001. [DOI] [PubMed] [Google Scholar]
- Clark C.E., Smith L.F.P., Taylor R.S., Campbell J.L. Nurse led interventions to improve control of blood pressure in people with hypertension: Systematic review and meta-analysis. *BMJ.* 2010;341:c3995. doi: 10.1136/bmj.c3995. [DOI] [PMC free article] [PubMed] [Google Scholar]
- Lambrinou E., Kalogirou F., Lammis D., Sourtzi P. Effectiveness of heart failure management programmes with nurse-led discharge planning in reducing re-admissions: A systematic review and meta-analysis. *Int. J. Nurs. Stud.* 2012;49:610–624. doi: 10.1016/j.ijnurstu.2011.11.002. [DOI] [PubMed] [Google Scholar]
- Moher D., Liberati A., Tetzlaff J., Altman D.G., The PRISMA Group Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement. *PLoS Med.* 2009;6:e1000097. doi: 10.1371/journal.pmed.1000097. [DOI] [PMC free article] [PubMed] [Google Scholar]

- Majed Haumed Al Zaidy¹, Sarah Talaq Almotairi², Afaf Abdullah Al-Dawsari³, Maha Othman Badri⁴, Lamia Nasser Alruwaili⁵, Latifa Rashid Almuteri⁶, Faizah Ateeg ALBoeinain⁷, Mona Hamdan Nahar Albelaasi⁵, Bashayer Ayed Farhan AlAnzi⁸, Alia Awad Farhan Al-Anzi⁸, Abdulrahman Bahri Madhan Al-Omran⁹, Ehsan sayeg alanzio
- Haynes R.B., Sacket D.L., Guyatt G.H., Tugwell P. *Clinical Epidemiology: How Clinical Practice Research*. 3rd ed. Lippincott Williams & Wilkins; Philadelphia, PA, USA: 2006. [Google Scholar]
- Higgins J.P.T., Altman D.G., Gøtzsche P.C., Jüni P., Moher D., Oxman A.D., Savovic J., Schulz K.F., Weeks L., Sterne J.A.C., et al. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ*. 2011;343:d5928. doi: 10.1136/bmj.d5928. [DOI] [PMC free article] [PubMed] [Google Scholar]
- Higgins J.P.T., Thompson S.G., Deeks J.J., Altman D.G. Measuring inconsistency in meta-analysis. *BMJ*. 2003;327:557. doi: 10.1136/bmj.327.7414.557. [DOI] [PMC free article] [PubMed] [Google Scholar]
- Song F., Khan K.S., Dinne J., Sutton A.J. Asymmetric funnel plots and publication bias in meta-analyses of diagnostic accuracy. *Int. J. Epidemiol.* 2002;31:88–95. doi: 10.1093/ije/31.1.88. [DOI] [PubMed] [Google Scholar]
- Krumholz H.M., Amatruda J., Smith G.L., Mattera J.A., Roumanis S.A., Radford M.J., Crombie P., Vaccarino V. Randomized trial of an education and support intervention to prevent readmission of patients with heart failure. *J. Am. Coll. Cardiol.* 2002;39:83–89. doi: 10.1016/S0735-1097(01)016990. [DOI] [PubMed] [Google Scholar]
- Koelling T.M., Johnson M.L., Cody R.J., Aaronson K.D. Discharge education improves clinical outcomes in patients with chronic heart failure. *Circulation*. 2005;111:179–185. doi: 10.1161/01.CIR.0000151811.53450.B8. [DOI] [PubMed] [Google Scholar]
- Aldamiz-Echevarría Iraurgi B., Muñoz J., Rodríguez-Fernández J.A., Vidán-Martínez L., Silva-César M., Lamelo-Alfonsín F., Díaz-Díaz J.L., Ramos-Polledo V., Castro-Beiras A. Randomized controlled clinical trial of a home care unit intervention to reduce readmission and death rates in patients discharged from hospital following admission for heart failure. *Rev. Esp. Cardiol.* 2007;60:914–922. doi: 10.1157/13109644. [DOI] [PubMed] [Google Scholar]
- Kommuri N.V.A., Johnson M.L., Koelling T.M. Relationship between improvements in heart failure patient disease specific knowledge and clinical events as part of a randomized controlled trial. *Patient Educ. Couns.* 2012;86:233–238. doi: 10.1016/j.pec.2011.05.019. [DOI] [PubMed] [Google Scholar]
- Cockayne S., Pattenden J., Worthy G., Richardson G., Lewin R. Nurse facilitated self-management support for people with heart failure and their family carers (SEMAPHFOR): A randomised controlled trial. *Int. Nurs. Stud.* 2014;5:1207–1213. doi: 10.1016/j.ijnurstu.2014.01.010. [DOI] [PubMed] [Google Scholar]
- De Souza E.N., Rohde L.E., Ruschel K.B., Mussi C.M., Beck-da-Silva L., Biolo A., Clausell N., RabeloSilva E.R. A nurse-based strategy reduces heart failure morbidity in patients admitted for acute decompensated heart failure in Brazil: The HELEN-II clinical trial. *Eur. J. Heart Fail.* 2014;16:1002–1008. doi: 10.1002/ejhf.125. [DOI] [PubMed] [Google Scholar]
- Boyde M., Peters R., New N., Hwang R., Ha T., Korczyk D. Self-care educational intervention to reduce hospitalisations in Heart Failure: A randomised controlled trial. *Eur. J. Cardiovasc. Nurs.* 2018;17:178–185. doi: 10.1177/1474515117727740. [DOI] [PubMed] [Google Scholar]
- Huynh Q.L., Whitmore K., Negishi K., Marwick T.H., ETHELRED Investigators Influence of risk on reduction of readmission and death by disease management programs in heart failure. *J. Card. Fail.* 2019;25:330–339. doi: 10.1016/j.cardfail.2019.01.015. [DOI] [PubMed] [Google Scholar]
- Albert N.M., Cohen B., Liu X., Best C.H., Aspinwall L., Prarr L. Hospital nurses' comfort in and frequency of delivering heart failure self-care education. *Eur. J. Cardiovasc. Nurs.* 2015;14:431–440. doi: 10.1177/1474515114540756. [DOI] [PubMed] [Google Scholar]
- Bramhall E. Effective communication skills in nursing practice. *Nurs. Stand.* 2014;29:53–59. doi: 10.7748/ns.29.14.53.e9355. [DOI] [PubMed] [Google Scholar]