JOURNAL OF INTERNATIONAL CRISIS AND RISK COMMUNICATION RESEARCH
ISSN: 2576-0017
2024, VOL 7, NO S12

Relation between Knowledge and Anxiety Level of Saudi Patients Undergoing Upper Gastrointestinal Endoscopy in 2024

Hanaa Mustafa Felemban[1], Jamilah Mustafa Felemban[2], Aqeel, Omair Munaji J[3], Hamlan, Duhaim Ibrahim M[3], Hattan, Wajdi Mohammed O[3], Khawaji, Dalia Ali I[3], Almash, Mohammed Amer A[3], Shawesh, Alhassan Hussain A[3], Almubarak, Suha Hussain A[3], Alhgwi, Najwa Mofarh A[3], Alhasani, Hamad Hussain A[3]

1 Midwife, Maternity and Children hospital, Saudi Arabia
2 Specialist Nursing Maternity and Children hospital, Saudi Arabia
3 NURSING, Jazan Health Cluster, Saudi Arabi.

Abstract*:* Upper Gastrointestinal (GI) endoscopies are essential for detecting and treating various digestive tract problems. While typically safe. Lack of patient's knowledge regarding endoscopy procedure is the main reason for increasing anxiety. ***This study aimed to:*** Assess knowledge and anxiety levels of patients undergoing upper gastrointestinal endoscopy. ***The study was carried out:*** In the gastrointestinal endoscopy units, Saudi university Hospital: A convenience sample of 150 adult patients with GI disorders admitted to GI endoscopy units and scheduled for GI endoscopy were recruit to the study. ***The research questions include***, 1- What are the levels of knowledge of patients undergoing upper GI endoscopy at Tanta University Hospital? 2- What are relations between levels of state and trait anxiety and different variables? ***Tools of the study:*** Two tools were used, **Tool (I) Structured interview questionnaire:** This tool was developed and used by the researcher for collection of baseline data and is consisted of three parts: Socio demographic data, patient's clinical information and patient's knowledge regarding upper gastrointestinal endoscopy. **Tool (II) The State Trait Anxiety Inventory Scale:** The original scale was developed by **Spielberger** in **1970** to evaluate respectively, state and trait anxiety. ***Results:*** It was observed that Less than half (44.00%) of studied patients had poor knowledge score, while (45.33%) had severe state anxiety and more than half (51.33%) had severe trait anxiety. there was a high negative significant correlation (r= -0.256, -0.318 respectively) between knowledge and The State Trait Anxiety Inventory Scale. ***Conclusion:*** Providing information to the patient before upper gastrointestinal endoscopy, decrease their level of anxiety and improve tolerance level during endoscopy procedure. ***Recommendations:*** It was recommended that provision of institutional written policies and guidelines regarding increasing knowledge and declining anxiety for patients undergoing upper gastrointestinal endoscopy.
Key words: Upper gastrointestinal endoscopy, knowledge, anxiety.

Introduction:

Endoscopy of the upper gastrointestinal tract (GI) is a common and safe treatment. Additionally, endoscopy has evolved beyond its use as a diagnostic tool to become a low-invasive, quick, and effective treatment technique. Despite the fact that improved endoscopic techniques result in higher-quality images, patient compliance is the primary factor influencing operation quality. The patient's degree of anxiety before to the treatment is one of the most significant elements influencing patient compliance (Al-Hagawi et al., 2024)

Globally, 750 gastroscopies surgery per 100.000 populations are subjected in 2004. Approximately 3.5 million people underwent gastroscopy in United States in 2016 (Hiremath et al., 2016). In Saudi Arabia, study done by Alruzug et al., 2021 found that a total of 2075 patients

were included with a mean age of 56.8 years (range 18–113) and males constituted 67.9%, while 65.9% had at least one comorbidity. Symptoms on presentation included hematemesis (52.5%), melena (31.2%), both hematemesis & melena (15.1%), and hematochezia (1.2%). The majority of UGIB were from a NVUGIB source (80.5%) and a variceal source was found in 13.1%, while no endoscopic findings were found in 6.4% of cases. The most common endoscopic diagnosis was gastroduodenal erosions (23.8%), duodenal ulcers (23.5%), reflux esophagitis (16.0%), esophageal varices (12.1%), and gastric ulcers (10.8%). There was no change in the endoscopic findings over the time period of the study. A third of duodenal ulcers (33.3%) as well as 21.9% of gastric ulcers were actively bleeding at the time of endoscopy, while 3.3% of duodenal ulcers had an adherent clot.

The esophagus and upper gastrointestinal tract (UGIT) are examined by upper gastrointestinal endoscopy. It is regarded as the preferred test for dyspepsia, reflux, or alarm symptoms, and it is necessary for an accurate diagnosis in these upper abdominal complaints. Upper gastrointestinal endoscopy procedure usually done for the purpose of diagnosing any upper GI problems and usually create anxiety for patient. Anxiety in patient occurs due to lack of awareness about the procedure and fear of discomfort or pain in procedure throughout (Rudin et al., 2007; Van et al, 2005)

Anxiety is a feeling of fear, uneasiness and worry. Usually generalized and unfocused, it is often accompanied by muscular tension, restlessness and problems in concentration. Upper gastrointestinal endoscopy nurses play a critical role in the provision of safe, high quality preparation or care for patients undergoing upper GI endoscopy such as preparing the endoscopic room with the right instrument and necessary devices for examination of the upper GIT (Gomez et al., 2009; Salvetto., 2007)

Moreover, the nurse offers a holistic package of care to patients undergoing upper GI endoscopy, encompassing the psychological, physiological and sociological needs of the patient. Also, the nurse already has the skills and knowledge to assess the needs of each individual attending for endoscopy from admission to discharge. In addition to, providing information to the patient before upper GI endoscopy may decrease the level of anxiety and improve tolerance level of patient during endoscopy procedure. To ensure the quality and safety of the procedure in upper GI endoscopy, it is important to know the patient anxiety level and the effective factors to reduce it, as enhance patient knowledge related to procedure (Koloski et al., 2009; El-Hassan et al., 2009; Addolorato et al., 2008; Tina et al., 2006; Pehlivan et al., 2011) .

Assessment is the first step of the nurse and patient interaction. It is also the most important aspect of the pre and post upper GI endoscopy procedure. In addition, to ensure individuality of care by finding out what the patient want and need to know. Also, help in clarify misconceptions and allow patient to express feeling. The pre procedure nursing assessment of the patient includes reviewing of medical history of the patient regarding allergy to any medicines, history of taking any medication, surgery and radiation treatments to esophagus, stomach or duodenum. Also, the nurse asks the patient taking nothing by mouth for about six hours, changing into a loose fitting gown, intravenous access be established, spectacles and dentures are removed and the informed consent sign by the patient and put in the patient record (Berzin et al., 2020).

The intra procedure nursing assessment includes frequent check and records all vital signs, and put the patient in appropriate position for performing the procedure usually the left side. A mouth guard place in the mouth to protect the patient teeth from the endoscopy. Then the lubricate tip of the endoscopy guide smoothly and slowly into the mouth and down into the esophagus, additionally the endoscopy nursing staff asks the patient to swallow to assist in moving the tube along. The post procedure nursing assessment includes frequent check and records all vital signs,

assesses patient's response to procedure as nausea, bloating, level of consciousness, ability to swallow and move extremities. If the patient's throat numb after the procedure, the patient doesn't eat or drink until throat is no longer numb, and gag reflex return to normal [(24)]. It is also crucial that, the nurse gives the right information before the procedure to relieve patient's anxiety. During the procedure the nurse must help the endoscopist. After the completion of the procedure, the nurse manages the endoscopic finding safely as biopsies (Bennett et al., 2016; Yoenlee et al., 2012) .

Subjects & Method:

Subjects:

The aim of this study is to assess knowledge and anxiety levels of patients undergoing upper GI endoscopy at Saudi University Hospital.

Research questions:

- What are the levels of knowledge of patients undergoing upper GI endoscopy at Saudi University Hospital?
- What are relations between levels of state and trait anxiety and different variables?

Study design:

Descriptive and exploratory study design was utilized to achieve the aim of the study and answer the research questions.

Setting of the study:

This study was conducted in the gastrointestinal endoscopy units at Saudi University Hospital.

Subjects:

A convenience sample of 150 adult patients with GI disorders admitted to GI endoscopy units at Saudi University Hospital and scheduled for GI endoscopy were recruit to the study. The subject size of participants was calculated using a power analysis.

The inclusion criteria were as follow:

- Ages ranged from 18 to 60 years.
- Both sexes.
- Conscious patients.
- First time undergoing upper GI endoscopy.

The exclusion criteria were as follow:

Patients taking anti-anxiety drugs in the last 72h.

- Patients unable to communicate.
- Patients with dementia or psychiatric disorders.
- Patient in emergency condition.

Data collection tools:

Two tools were used in this study to collect pertinent data related to the aim as follow:

Tool (I): Structured Interview Questionnaire:

This tool was developed by the researcher after review of the relevant literature to assess patient's socio demographic data, clinical information and their knowledge about upper GI endoscopy. It consisted of three parts as follow:

Part 1: Socio demographic data:

It was developed to assess patient's socio demographic data that covered the following variables: age, sex, marital status, place of residence, socioeconomic status, level of education, income level and occupation.

Part 2: **Patient's clinical information:** This part was consisted of statements that used to assess patients information about their health history, it was comprise the following areas: present diagnosis, past medical history, history of hospitalization, the last laboratory studies, heart rate, respiration rate, blood pressure, associated chronic diseases, mobility status, past and current intake of medication, the last attack of bleeding, previous time of bleeding factors that participate in bleeding such as certain drug, smoking, lifting heavy objects, constipation, coughing or sneezing (Bennett et al., 2016; Yoenlee et al., 2012).

Part 3: Patient's knowledge assessment:

This part was developed to assess patient's knowledge about upper GI endoscopy such as: Definition, indication, purposes, risk factors, procedure, possible occurrence of discomfort or side effect, the place where the procedure was performed, investigation was done, type of anesthesia used, physical and psychological preparation before procedure. Also, what was actually happen during the procedure, recovery period and discharge instruction such as follow up plan, types of activities, return to work, types of diet, medication schedule, hygiene care, signs of complication and referral places (Adang et al., 2010).

Scoring system of patient's knowledge assessment questionnaire was done as follow:

Patient's knowledge assessment questionnaire was consisted of 22 statements about knowledge and information of the patients regarding upper GI endoscopy. Each question has a number of a group of answer points.

Scoring system: Patient who was responded by correct and complete answer was given a score two, correct and incomplete answer was given a score one and the patients who responded wrong and not answer was given a score zero. Less than 60% of total score was considered as poor, from 60% to less than 75% was considered as fair, and from75% and more was considered as good.

Tool (II): The State Trait Anxiety Inventory Scale (STAI):

The original scale the State Trait Anxiety Inventory was developed by **Spielberger** in **1970**[28]. It was used to evaluate respectively, state and trait anxiety. The SAI Scale determines how the individual feels at a particular moment and under certain circumstances. The scale is designed in quadruple Likert-type scale varying between never, sometimes, usually and almost always. The TAI Scale determines how the individual generally feels, independent of the status and the conditions which surround him/her (Elsayd et al., 2016).

Scoring system: Both the SAI and TAI Scales comprised of 20 items each one scored on 4 point forced choice Likert type response scales. Scores ranged from 20 to 80, with normal scores or no anxiety score 20, low scores from 21-40 suggest mild anxiety, median scores from 41-60 suggest moderate anxiety, while high scores from 61-80 suggest severe anxiety.

Method:

Approval was obtained from the responsible authority of GI endoscopy units at Saudi University Hospitals before conducting the study.

Written consent was obtained from every patient included in the study after explanation of the aim of the study and assuring them of confidentiality of collected data. Confidentiality and anonymity was maintained by the use of code number instead of name and the right of withdrawal is reserved.

Privacy was assured to the patients.

Tools development: Tools I (Structured Interview Questionnaire) was developed by researcher based on relevant literature review for collection of baseline data

Tool II (The STAI Scale) was developed by **Spielberger** in **1970**, and were initially designed as a single instrument to measure both anxiety trait and anxiety state. Arabic translation of this tool was done by **El-Behairy** in **1984** to a certain relevance and completeness.
All tools were tested for content validity by a nine jury of experts in the field and Modifications were done to certain relevance and completeness.

All tools of the study were tested for reliability and Cronbach alpha was used and found to be 0.88 for Tool I and 0.98 for Tool II, which consider highly reliable tools.

A pilot study was carried out on a sample of 150 patients from GI endoscopy unit in order to test clarity, feasibility and applicability of the tools. Subject of pilot study was excluded from the original sample.

Data collection was conducted over a period of 6 months (started from first of January to the end of June 2024). After data collection, data was coded, analyzed then tabulated under the direction of a statistician to obtain results to answer the research questions.
Finally, most patients approach an endoscopic procedure with fear and anxiety. Moreover, to lessen or even prevent this, providing patients with information is essential in order to prepare the patients physically, emotionally and intellectually for the procedure of upper GI endoscopy.

Statistical analysis:

After completion of data collection, each sheet of interview questionnaire sheet information and STAI Scale questions was coded, organized and categorized then the data was tabulated and presented into frequency distribution tables. The following tests used in the study were chi square test and P-value to assess knowledge and anxiety levels of patients undergoing upper GI endoscopy at Tanta University Hospital. The data was collected and statistically analyzed using the Statistical Package for Social Sciences (SPSS) version 21 for continuous variables (mean ±SD), ANOVA t tests were used for the detection of significant differences for studied patients. Pearson's correlation coefficient was used to test correlation between variables.
10. The level of significance chose in the study was set at 0.05 levels.
Non significance if P-value > 0.05
Significance if P-value < 0.05
High significance if P-value < 0.001

Results:

Table (1) illustrates percentage distribution of studied patients according to socio demographic characteristics. This table showed that the mean age of studied patients was (52.56±12.07). More than half (60.67%) in the age group ranged from 51 to 60 years old, while (15.33%) in the age group ranged from 30 to 40 years old. **In relation to sex**, more than two thirds (71.33%) of studied patients were male, while less than one third (28.67%) were female. **As regards to marital status,** more than three quarters (82.67%) of studied patients were married, while (4.00%) were divorced. **In relation to residence,** less than three fourths (72.00%) of studied patients have lived in rural areas, while less than one third (28.00%) have lived in urban areas.
As regards to income level, less than two thirds (62.67%) of studied patients had low income level, while (12.00%) had high income level. **In relation to level of education,** more than one

half (54.00%) of studied patients were illiterate, while (6.67%) had secondary education.
As regards to occupation, less than two thirds (64.00%) of studied patients had no work, while more than one third (36.00%) working.

Variables	n (150)	%
Age (years)		
-30-40	23	15.33
-41-50	36	24.00
-51- 60	91	60.67
-Mean±SD	52.56±12.07	
Sex		
-Male	107	71.33
-Female	43	28.67
Marital status		
-Single	10	6.67
-Married	124	82.67
-Divorced	6	4.00
-Widow	10	6.67
Residence		
-Urban	42	28.00
-Rural	108	72.00
Income level		
-Low	94	62.67
-Middle	38	25.33
-High	18	12.00
Education		
-Illiterate	81	54.00
-Read and write	31	20.67
-Primary education	12	8.00
-Secondary education	10	6.67
-University	16	10.67
Occupation		
-Work	54	36.00
-Not work	96	64.00

Table (2) shows distribution of studied patients according to current diagnosis and past medical history. This table illustrated that less than half (44.67%) of studied patients referred from other department to endoscopy units, while, (37.33%) and (18.00%) of studied patients were admitted from outpatients and emergency departments respectively. **In relation to current diagnosis,** It can be seen that the most common diagnosis of studied patients who undergoing upper GI endoscopy was esophageal varices and ulcer (28.67%), while the least diagnosis was abdominal pain and distention (5.33%). **As regards to past medical history,** more than one third (36.00%) of studied patients had past medical history of liver and spleen diseases, while (2.00%) had past medical history of cancer. **Finally, regarding history of hospitalization,** more than half (54.67%) of studied patients had no previous history of hospitalization, while less than half (45.33%) had previous history of hospitalization and nearly two thirds (63.24%)

hospitalized for medical reasons.

Variables	n	%
Admitted from		
-Outpatients	56	37.33
-Emergency	27	18.00
-Referred *	67	44.67
Diagnosis		
-Esophageal varices and ulcer	43	28.67
Esophageal varices grade I-	18	12.00
-Esophageal varices grade II	33	22.00
-Esophageal varices grade III	18	12.00
-Esophageal stenosis	12	8.00
-Gastric ulcer	18	12.00
-Abdominal pain and distention	8	5.33
Past medical history		
-No	38	25.33
-Diabetes mellitus	23	15.33
-Renal diseases	5	3.33
-Hypertension	11	7.33
-Liver and spleen diseases	54	36.00
Pulmonary diseases-	5	3.33
-Cancer	3	2.00
Heart diseases-	6	4.00
Blood diseases-	5	3.33
Family history		
-No	113	75.33
-Diabetes mellitus	16	10.67
-Renal diseases	7	4.67
-Liver diseases	11	7.33
-Heart diseases	1	0.67
-Blood diseases	2	1.33
History of hospitalization		
Yes-	68	45.33
-No	82	54.67
If yes, for what		
-Medical**	43	63.24
-Surgical	25	36.76

Table (3) illustrates distribution of studied patients according to history of upper gastrointestinal bleeding. One can notice that, more than half (57.03%) of studied patients had information about upper GI endoscopy from medical team, while (28.13% and 23.44%) of studied patients had their information from friends and media respectively. As regards to factors that participate **bleeding** less than two thirds (62.07%) of studied patients had no factors participate in bleeding, while (37.93%) had factors that participate bleeding. The most common precipitating factors of bleeding was coughing or sneezing (36.36%), while the least one was constipation (15.91%).

Variables	n	%
Source of information about endoscopy*		
-Medical team	73	57.03
-Friends	36	28.13
-Media	30	23.44
History of upper gastrointestinal (GI) bleeding		
-Yes	116	77.33
-No	34	22.67
	n=116	
If yes, how many times		
-One	59	50.86
-Two	26	22.41
-More	31	26.72
The last attack of bleeding		
-< one month	49	42.24
-1- < 6 months	8	6.90
-6 months - one year	31	26.72
-> one year	28	24.14
There are factors that participate bleeding		
-Yes	44	37.93
-No	72	62.07
	n=44	
If yes, what are these		
-Constipation	7	15.91
-Coughing or sneezing	16	36.36
-High blood pressure	9	20.45
-Emotional stress	12	27.27

Figure (1): Percentage distribution of studied patients in relation to their total knowledge score regarding upper gastrointestinal endoscopy.

This figure showed that less than half (44.00%) of studied patients had poor knowledge score regarding upper GI endoscopy, while (24.00% and 32.00%) of studied patients had fair and good knowledge score respectively.



Table (4) illustrates percentage distribution of the studied patients in relation to their total Anxiety Inventory Scale pre upper gastrointestinal endoscopy procedure. It can be seen that less than half (45.33%) of studied patients had severe state anxiety, while (18.67%, 10.00% and 26.00%) had normal, low and moderate state anxiety respectively. **Regarding total Anxiety Trait Scale,** more than half (51.33%) of studied patients had severe trait anxiety, while (12.00%, 16.00% and 20.67%) had normal, low and moderate trait anxiety respectively. It was found that, highly statistical significant difference in relation to patients total anxiety state and total anxiety trait pre upper GI endoscopy procedure**.**

	Anxiety Inventory Scale									
	Normal		**Low**		**Moderate**		**Severe**		**Chi-square**	
	n	**%**	**N**	**%**	**n**	**%**	**n**	**%**	**X^2**	**P-value**
-The State	28	18.67	15	10.00	39	26.00	68	45.33	40.773	<0.001**
-The Trait	18	12.00	24	16.00	31	20.67	77	51.33	57.733	<0.001**

Table (5) reveals relation between patient's total knowledge score and The State Anxiety Inventory Scale. It can be seen that, (50.00%) of studied patients had good knowledge and normal state anxiety scale, while more than half (58.82%) of studied patients had poor knowledge and severe state anxiety scale. **Regarding relation between total knowledge score and The Trait Anxiety Scale,** half (50.00%) of studied patients had good knowledge and normal trait anxiety scale, while more than half (59.74%) of studied patients had poor knowledge and severe trait anxiety scale. It can be seen that, there was a statistical significant difference among studied patients in relation to knowledge and The Anxiety Inventory Scale, p-value was <0.05.

Anxiety Inventory Scale	**Knowledge Score**									
	Poor		**Fair**		**Good**		**Total**		**Chi-square**	
	n	**%**	**N**	**%**	**n**	**%**	**n**	**%**	**X^2**	**P-value**
The State							**The State (r=-0. 256/P=0.020*)**			

-Normal	5	17.86	9	32.14	14	50.00	**28**	**18.67**	15.466	0.017*
-Low	7	46.67	4	26.67	4	26.67	**15**	**10.00**		
-Moderate	14	35.90	10	25.64	15	38.46	**39**	**26.00**		
-Severe	40	58.82	13	19.12	15	22.06	**68**	**45.33**		
The Trait							**The Trait (r=-0. 318/ P<0.001**)**			
-Normal	3	16.67	6	33.33	9	50.00	**18**	**12.00**	23.510	<0.001**
-Low	9	37.50	6	25.00	9	37.50	**24**	**16.00**		
-Moderate	8	25.81	6	19.35	17	54.84	**31**	**20.67**		
-Severe	46	59.74	18	23.38	13	16.88	**77**	**51.33**		

Table (6) reveals relation between socio demographic data of patients and their total knowledge score. This table showed that the mean age of studied patients who had good knowledge was (52.18±8.35). **In relation to sex**, less than one third (31.33%) of studied patients who had poor knowledge were male, while (12.67%) were female. On other hands, more than one fifth (20.67%) of studied patients who had good knowledge were male. **As regards to marital status,** nearly one third (32.67%) of studied patients who had poor knowledge were married. While, (26.67%) who had good knowledge were married. **In relation to residence,** more than one third (34.67%) of studied patients who came from rural had poor knowledge, compared to (9.33%) came from urban. While, (21.33%) of studied patients who came from rural had good knowledge. **In relation to level of education,** more than one quarter (25.33%) of studied patients who had poor knowledge were illiterate, while (2.00%) were graduated from university. On the other hand, (13.33%) of studied patients who had good knowledge were illiterate, while (8.00%) were graduated from university. It can be seen that, there was a highly statistical significant difference between knowledge in relation to sex, marital status, socioeconomic status, income level and occupation, p-value was <0.05*

Variables	**Knowledge**									
	Poor		**Fair**		**Good**		**Total**		**Tests**	
	n	**%**	**n**	**%**	**n**	**%**	**n**	**%**	**X^2 or f**	**P-value**
Age										
-Mean±SD	47.32±11.15		49.67±10.62		52.18±8.35				3.695	0.027*
Sex										
-Male	47	31.33	29	19.33	31	20.67	**107**	**71.33**	2.567	0.277
-Female	19	12.67	7	4.67	17	11.33	**43**	**28.67**		
Marital status										
-Single	6	4.00	1	0.67	3	2.00	**10**	**6.67**	23.030	<0.001**
-Married	49	32.67	35	23.33	40	26.67	**124**	**82.67**		
-Divorced	1	0.67	0	0.00	5	3.33	**6**	**4.00**		
-Widow	10	6.67	0	0.00	0	0.00	**10**	**6.67**		
Residence										
-Urban	14	9.33	12	8.00	16	10.67	**42**	**28.00**	2.694	0.260

-Rural	52	34.67	24	16.00	32	21.33	**108**	**72.00**		
Education										
-Illiterate	38	25.33	23	15.33	20	13.33	**81**	**54.00**	23.205	0.003*
-Readand write	12	8.00	9	6.00	10	6.67	**31**	**20.67**		
-Primary edu	9	6.00	2	1.33	1	0.67	**12**	**8.00**		
-Secondary edu	4	2.67	1	0.67	5	3.33	**10**	**6.67**		
-University	3	2.00	1	0.67	12	8.00	**16**	**10.67**		
Income level										
-Low	50	33.33	23	15.33	21	14.00	**94**	**62.67**	23.745	<0.001**
-Middle	13	8.67	12	8.00	13	8.67	**38**	**25.33**		
-High	3	2.00	1	0.67	14	9.33	**18**	**12.00**		
Occupation										
-Work	8	5.33	12	8.00	34	22.67	**54**	**36.00**	41.723	<0.001**
-Not work	58	38.67	24	16.00	14	9.33	**96**	**64.00**		

Table (7) Illustrates relation between socio demographic data of patients and The State Anxiety Inventory Scale.

This table showed that the mean age of studied patients who had severe state anxiety was (51.67±8.51). **In relation to sex**, less than one third (28.67%) of studied patients who had severe state anxiety were male, while (16.67%) were female. On other hand, (7.33%) of studied patients who had low state anxiety were male, while (2.67%) were female. **As regards to marital status,** more than one third (36.67%) of studied patients who had severe state anxiety were married, while (2.67%) were single. On other hand, (8.00%) of studied patients who had low state anxiety were married, while (2.00%) were single. **In relation to residence,** the percentage (33.33% and 12.00%) of studied patients who came from rural and urban areas had severe state anxiety respectively, while (6.00% and 4.00%) of studied patients who came from rural and urban had low state anxiety respectively. **In relation to level of education,** more than one fifth (23.33%) of studied patients who had severe state anxiety were illiterate, while (2.67%) were graduated from university. Also, (6.00%) of studied patients who had low state anxiety were illiterate. It can be seen that, there was a highly statistical significant difference between the SAI Scale in relation to age, marital status, socioeconomic status and education level, p value was <0.05*

Variables	**The State Anxiety Inventory scale**											
	Normal		**Low**		**Moderate**		**Severe**		**Total**		**Tests**	
	n	**%**	**n**	**%**	**n**	**%**	**n**	**%**	**n**	**%**	**X^2 or f**	**P-value**
Age (years) -Mean±SD	56.29±11.48		54.26±12.19		50.17±10.24		51.67±8.51				2.373	0.073
Sex												
-Male	24	16.00	11	7.33	29	19.33	43	28.67	**107**	**71.33**	5.216	0.157
-Female	4	2.67	4	2.67	10	6.67	25	16.67	**43**	**28.67**		

Marital status												
-Single	1	0.67	3	2.00	2	1.33	4	2.67	**10**	**6.67**	15.316	0.083
-Married	24	16.00	12	8.00	33	22.00	55	36.67	**124**	**82.67**		
-Divorced	3	2.00	0	0.00	2	1.33	1	0.67	**6**	**4.00**		
-Widow	0	0.00	0	0.00	2	1.33	8	5.33	**10**	**6.67**		
Residence												
-Urban	4	2.67	6	4.00	14	9.33	18	12.00	**42**	**28.00**	4.969	0.174
-Rural	24	16.00	9	6.00	25	16.67	50	33.33	**108**	**72.00**		
Education level												
-Illiterate	14	9.33	9	6.00	23	15.33	35	23.33	**81**	**54.00**	23.921	0.021*
-Read and write	3	2.00	3	2.00	9	6.00	16	10.67	**31**	**20.67**		
-Primary edu	0	0.00	0	0.00	1	0.67	11	7.33	**12**	**8.00**		
-Secondary edu	4	2.67	1	0.67	3	2.00	2	1.33	**10**	**6.67**		
-University	7	4.67	2	1.33	3	2.00	4	2.67	**16**	**10.67**		
Income level												
-Low	14	9.33	7	4.67	24	16.00	49	32.67	**94**	**62.67**	11.806	0.066
-Middle	7	4.67	6	4.00	13	8.67	12	8.00	**38**	**25.33**		
-High	7	4.67	2	1.33	2	1.33	7	4.67	**18**	**12.00**		
Occupation												
-Work	15	10.00	6	4.00	17	11.33	16	10.67	**54**	**36.00**	9.421	0.024*
-Not work	13	8.67	9	6.00	22	14.67	52	34.67	**96**	**64.00**		

Table (8) reveals relation between socio demographic data of patients and The Trait Anxiety Inventory Scale. This table showed that the mean age of studied patients who had severe trait anxiety was (51.62±11.58). **In relation to sex**, more than one third (35.33%) of studied patients who had severe trait anxiety were male, while (16.00%) were female. On the other hand, (12.00%) of studied patients who had low trait anxiety were male, while (4.00%) were female. **As regards to marital status,** more than two fifth (40.67%) of studied patients who had severe trait anxiety were married, while (2.67%) were single. On other hands (12.67%) of studied patients who had low trait anxiety were married, while (2.67%) were single. **Regarding level of education,** (30.00%) of studied patients who had severe trait anxiety were illiterate, compared to (4.00%) of them were graduated from university. While (8.67%) of studied patients who had low trait anxiety were illiterate, compared to (2.00%) of them were graduated from university.

Variables	**The Trait Anxiety Inventory scale**										**Tests**	
	Normal		**Low**		**Moderate**		**Severe**		**Total**			
	n	**%**	**N**	**%**	**n**	**%**	**n**	**%**	**n**	**%**	**X^2 or f**	**P-value**
Age (years) -Mean±SD	53.66±12.82		55.92±8.7		52.73±12.54		51.62±11.58				0.887	0.450
Sex												

-Male	14	9.33	18	12.00	22	14.67	53	35.33	**107**	**71.33**	0.761	0.859
-Female	4	2.67	6	4.00	9	6.00	24	16.00	**43**	**28.67**		
Marital status											9.851	0.363
-Single	1	0.67	4	2.67	1	0.67	4	2.67	**10**	**6.67**		
-Married	16	10.67	19	12.67	28	18.67	61	40.67	**124**	**82.67**		
-Divorced	1	0.67	0	0.00	1	0.67	4	2.67	**6**	**4.00**		
-Widow	0	0.00	1	0.67	1	0.67	8	5.33	**10**	**6.67**		
Residence											8.805	0.032*
-Urban	7	4.67	11	7.33	10	6.67	14	9.33	**42**	**28.00**		
-Rural	11	7.33	13	8.67	21	14.00	63	42.00	**108**	**72.00**		
Education											15.658	0.207
-Illiterate	8	5.33	13	8.67	15	10.00	45	30.00	**81**	**54.00**		
-Read and write	2	1.33	7	4.67	6	4.00	16	10.67	**31**	**20.67**		
-Primary education	1	0.67	0	0.00	3	2.00	8	5.33	**12**	**8.00**		
-Secondary education	4	2.67	1	0.67	3	2.00	2	1.33	**10**	**6.67**		
-University	3	2.00	3	2.00	4	2.67	6	4.00	**16**	**10.67**		
Income level											8.239	0.221
-Low	7	4.67	16	10.67	18	12.00	53	35.33	**94**	**62.67**		
-Middle	6	4.00	6	4.00	10	6.67	16	10.67	**38**	**25.33**		
-High	5	3.33	2	1.33	3	2.00	8	5.33	**18**	**12.00**		
Occupation											18.816	<0.001**
-Work	13	8.67	9	6.00	15	10.00	17	11.33	**54**	**36.00**		
-Not work	5	3.33	15	10.00	16	10.67	60	40.00	**96**	**64.00**		

Table (9) reveals correlation between studied patient's total knowledge score **and The State Trait Anxiety Inventory Scale.** It can be seen that, there was a high negative significant correlation (r = - 0.256, -0.318 respectively) between knowledge and The STAI Scale, P value<0.001. On other hands, there was a high positive significant correlation (r=0.712) between The SAI Scale and The TAI Scale, P value <0.001.

Variables	**Knowledge**		**The STAI Scale**	
	R	**P-value**	**r**	**P-value**
The State Anxiety Inventory (SAI) Scale	-0.256	0.020*		
The Trait Anxiety Inventory (TAI) Scale	-0.318	<0.001**	0.712	<0.001**

Discussion:

The upper gastrointestinal endoscopic procedure anxiety was often associated with the fear of experiencing pain and discomfort during endoscopy, decreases patient cooperation and tolerance

with procedure. Therefore, control of discomfort and pain before the endoscopy was considered to be a high priority by patients. The best intervention to reduce anxiety is to inform patient with adequate knowledge about the procedure (Sewitch et al., 2011). For these reasons, the current study aimed to assess knowledge and anxiety level of patients undergoing upper GI endoscopy

Less than two thirds of studied patients who undergoing upper GI endoscopy was in age group ranged from 51 to 60 years. This could be explained that this age represents last period of working age population which exposed to unhealthy life style such as stressful life events, misuse of medications as analgesics that effect on gastro intestinal tract, and increased consumption of fast food, spicy and high fat diet. This finding was in agreement with **Ruhl and Everhart (2016)**who stated that the majority of patients undergoing upper GI endoscopy belonged to the age group 50-59 years.
On other hand, this finding was contradicted by **Lee et al (2012)** and **El said (2016)**who mentioned that, most of patients undergoing upper GI endoscopy were belonged to the age 41years.
Moreover, in the current study it was found that more than two thirds of studied patients were male. This could be explained that male patients exposed to stressful life, heavy smoking and increasing consumption of caffeine more than female patients. This finding was highly supported with study done by **Seda et al (2011)** who found that the majority of studied patients who undergoing upper GI endoscopy were male.

Also this finding was inconsistent with **Smitha and Sugirtha** (**2015)**who stated that majority of patients were predominantly male. On the other hand, this result was not accordance with study done by **Feyzullah et al (2010)**who reported that upper GI endoscopy more common in women than in men.

Regarding marital status, the current study revealed that more than three quarters of studied patients were married, this finding was in harmony with **Mohammed(2016)** who recorded that nearly two thirds of studied patients were married. Also, this finding was in line with **Abdulla (2010)** who reported that the majority of patients in a study entitled; "Assessment of knowledge and perception of the patient with upper gastrointestinal disorders" were married.
As regards to residence, the current study showed that more than two thirds of studied patients had lived in rural. It may be due to people lived in rural area usually ignore any pain suffers and neglect routine checkup, also they doesn't find big hospitals provide intensive care. This finding inagreement with **Prabhuswami et al (2016)**who stated that the majority of studied patients came from rural area. Also, this result was supported by **Kennedy et al (2006** and **Vliet et al (2004**, they illustrated that rural patients with lowest socioeconomic status were high risk for upper GI diseases.

Concerning income level, in the current study it was found that less than two thirds of studied patients consider low income level, more than half were illiterate and nearly two thirds of them had no work. This may be interpreted that poor income and low educational levels had negative effects on health, and these events may affect knowledge and anxiety level among patients. This finding was supported by **Quick (2006)** who revealed that illiterate and low income level patient did not seek the hospital or clinics except in cases of necessity only, and for this reason the medical problem became severe.
In addition, this finding was consistent with **Winslow (2004)** and **Mohamed et al (2014)**, they mentioned that; the majority of studied patient who undergoing upper GI endoscopy were illiterate and had low income level. n the other hand, this finding was contradicted by study done with **Previti et al (2016)** who revealed that more than two thirds of studied patients who undergoing

upper GI endoscopy had high education level.

Upper gastrointestinal endoscopy procedure is a difficult, stressful, anxiety associated, unpleasant diagnostic and therapeutic method. Providing adequate information before upper GI endoscopy lead to decrease anxiety, fear and worry of the patients during the procedure. For that, an effective health teaching guidelines should be given, depending on the patient's individual needs and the patient should be assisted in order to establish good compliance throughout the procedure (Axon et al., 2010; Chair et al., 2005) .

Regarding patient's knowledge before upper GI endoscopy; it can be seen from current study that less than half of studied patients had unsatisfied knowledge score regarding upper GI endoscopy, while less than one third of them had good knowledge score .

Additionally, these results suggested that providing verbal information to the patients is effective and essential in increasing knowledge about the nature and duration of the upper GI endoscopy, helping them to feel secure and reducing their anxiety associated with the result of the procedure. This finding was contradicted by **Prabhuswami et al (2016)**, they mentioned that the majority of studied patients who undergoing upper GI endoscopy had good knowledge.

In order to provide sufficient information to studied patients before undergoing upperGI endoscopy, the information should cover different issues such as: Indication, types, possible occurrence of discomfort or side effect, physical and psychological preparation before endoscopy.Also, these findings were inagreement with study done by **Thomas and Sugirtha (2015)**, they illustrated that good preparation by providing adequate knowledge to patient before diagnostic GI endoscopy enhance patients understanding and awareness about this procedure.

The finding of this study showed that less than half of studied patients had severe state anxiety, while more than half had severe trait anxiety. These findings may due to lack of awareness about the upper GI endoscopy procedure, fear of discomfort or pain in procedure and/or its results. These findings were in similar with study done by **Jones et al(2004),** they reported that the anxiety level was increased before the endoscopy procedure.

Also, these findings were inline with **Brandt (2001)** who demonstrated that patients scheduled for endoscopy procedure were usually anxious from the procedure. On other hand, these findings were contradicted by **Trevisani et al (2016)** and **Essink-Bot et al (2016**, they revealed that no correlation between upper GI endoscopy and knowledge and pre procedural anxiety.

A statistical significant difference was observed among studied patients in relation to their knowledge and The STAI Scale due to structured information might decrease patient anxiety and improve general health. This result was supported with the study conducted by **Ajee (2002)and Aabakken (2010**, these studies emphasized that structured information decrease anxiety and improve patient's tolerance level.

In addition, similar results were found by **Kiyohara et al**(**2004)**who revealed that increase in patient's knowledge regarding upper GI endoscopy procedure reduced their anxiety level.

Moreover, it can be seen that nearly half of studied patients had good knowledge and normal STAI Scale, while more than half had poor knowledge and severe STAI Scale. This means that adequate information and increase awareness among patients pre upper GI endoscopy procedure associated with reduce the anxiety level. This finding was in agreement with study done by **Rohde** (**2011** who experienced severe anxiety before GI endoscopy had poor knowledge regarding the procedure.

Additionally, this finding was in line with the study done by **Peter et al (2008)**who showed that a combination of information and training prior to upper GI endoscopy procedure is an effective

means for reducing anxiety.

It was found that a high statistical significant difference between knowledge in relation to sex, marital status, socioeconomic status, education level, income level and occupation. It may be due to more than half of studied patients had low level of education, no occupation and poor socioeconomic status which reflect on acquisition of knowledge. However, those patients had the confidence in themselves and cognitive maturity to get the information that they need through medical teams, reading articles and discussion with others. This result was inconsistent with **Mulcahy et al (2010**, who demonstrated that besides anxiety, the age, education level and gender also influenced significantly the tolerance of studied patients to upper GI endoscopy.
On other hand, this result was contradicted by **Hashimot (2002)** who pointed that there was no significant difference in relation to sex or age with tolerance to endoscopy.

Also, **Lee et al (2004** and **Campo(2015**, they reported that no significant difference in relation to patient's age, gender and education level related to upper GI endoscopy.
The current study revealed that there was a high statistical significant difference found between The STAI Scale in relation to sex, in fact nearly two thirds of studied patients who had severe STAI were male, while nearly one third of them were female. It may be due to no equality number in both gender of studied sample in this study.
These findings contradicted by **Trevisani et al (2002**, **Muzzarelli et al (2006)** and **Hakan et al (2012**, they identified that anxiety prior to the upper GI endoscopy had been reported to be more severe in women due to fear of not waking up after the procedure than in men.

The present study demonstrated that there was a negative significant correlation between knowledge and The STAI Scale. It may be due to the recommendation of upper GI endoscopy procedure, increases the patient's apprehension and anxiety that is due to lack of knowledge of the patient about endoscopy, also awaiting the result of endoscopy. Moreover, most of studied patients were illiterate and consider low socioeconomic standard, so financial resource decreased, which lead to increase level of stress and anxiety (Ersoz et al., 2010).
On other hands, there was a positive significant correlation between The SAI and TAI. This means the internal personality, generally feel and trait in life of studied patients had positive effect on them in different situations as immediately before the procedure. This finding was consistent with study done by **Trevisani et al (2002)**, they found no statistical significant difference between The SAI and TAI in studied patients who undergoing upper GI endoscopy.

Conclusion and recommendations:

1. Conclusion

Based on the findings of the present study, it can be concluded that:

- There was a negative statistical significant relationship existed between knowledge and the anxiety scale of studied patients, the results revealed that studied patients with good knowledge score appeared in normal anxiety scale, while studied patients who had poor knowledge appeared in severe anxiety scale.

- Moreover, a positive statistical significant relationship was observed between SAI and TAI scales. In the present study, less than half of studied patients had severe SAI Scale, while more than half had severe TAI Scale.
- The study also revealed that, there were certain factors that influence the knowledge and The STAI Scale of studied patients as sex, marital status, socioeconomic status, income level and occupation.

- Finally, overall findings revealed that good preparation by providing information to the patient before the procedure, decrease their level of anxiety and improve tolerance level.

Recommendations

Based upon the findings of this study, the following recommendations are derived and suggested:

- Counseling should be provided for all patients who are undergoing upper GI endoscopy that helps in preparation of them and reducing anxiety level.
- Assessment of patient's knowledge towards upper GI endoscopy must be done upon patient admission by nurses.
- Assessment of patient's anxiety towards upper GI endoscopy must be done in the initial data collection and be documented in patients file by nurses using.

Reference:

- Aabakken L. Development and evaluation of written patient information for endoscopic procedure, Endoscopy J. 2010; 29: 23-6.
- Abdulla W. Assessment of knowledge and perception of the patient with upper gastrointestinal disease. Master Thesis, Faculty of Nursing, Ain Shams University. 2010.
- Adang R, Vismans J and Talmon J. Appropriateness of indications for upper gastrointestinal endoscopy: Associations with relevant endoscopic disease, Gastrointest Endosc J. 2010; 42:390-7.
- Addolorato G, Mirijello A and D'AngeloC. State and trait anxiety and depression in patients affected by gastrointestinal diseases: Psychometric evaluation of 1641 patients referred to an internal medicine outpatient setting, Int J Clin Pract. 2008; 62: 1063-9.
- Ajee K. A study to assess the effectiveness of structured teaching on anxiety and physiological responses of adult patients undergoing endoscopy procedure. MSc Nursing. College of Nursing CMC Vellore: Tamilnadu University 2002.
- Al-Hagawi, Y., Alqahtani, N. I., Alsharif, S. N., Chakik, R., Asiri, D. H., Al mani Sr, S. Y., ... & Abdullah, A. A. B. (2024). Endoscopy of Low BMI Patients Compared to Normal BMI Patients: A Study From the Aseer Region, Saudi Arabia. *Cureus*, *16*(5).
- Alruzug, I. M., Aldarsouny, T. A., Semaan, T., Aldaher, M. K., AlMustafa, A., Azzam, N., ... & Almadi, M. A. (2021). Time trends of causes of upper gastrointestinal bleeding and endoscopic findings. *Saudi Journal of Gastroenterology*, *27*(1), 28-34.
- Axon A, Jones R and Quine M. Guidelines on appropriate indications for upper gastrointestinal endoscopy, American Nursing J. 2010; 313 (5): 922-8.
- Bennett S, Gostimir M and Shorr R. The role of routine postoperative upper endoscopy in bariatric surgery, Surg Obes Relat Dis J. 2016; 10:755-65.
- Berzin T, Blanco P, Lamont J and Sawhney M. Persistent psychological or physical symptoms following endoscopic procedures: An unrecognized post endoscopy adverse event, Dig Dis Sci J. 2010; 55: 2869-73.
- Brandt L. Patients' attitudes and apprehensions about endoscopy: How to calm troubled waters. Americian Gastroenterol J. 2001; 96 (1): 280-4.
- Campo R, Brullet E and Montserrat A. Identification of factors that influence tolerance of upper gastrointestinal endoscopy, Eur Gastroenterol HepatolJ. 2015; 21:210-24.
- Chair S and Thompson D. Patient teaching prior to coronary angiography in Hong Kong: A pilot

study, Clinical Nursing J. 2005; 14(10): 114-5.

- El said S. Effectiveness of structured teaching program on knowledge, anxiety state and tolerance for patients with gastrointestinal endoscopy: Randomized controlled trial, Life Science Journal. 2016; 13(9): 9-17.
- El-Hassan H, McKeown K and Muller Music reduces anxiety levels in patients attending for endoscopy, Aliment Pharmacol Ther J. 2009;30(7):718–24.
- Elsayd O, Elshemy M and Elsays H. Effect of a multi-modal preparation package on knowledge and anxiety among patients undergoing cardiac catheterization, International Journal of Nursing Didactics. 2016; 6:1-10.
- Ersoz F, Toros A and Aydogan G. Assessment of anxiety levels in patients during elective upper gastro- intestinal endoscopy and colonoscopy, Turk Gastroenterol J. 2010; 21: 29-33
- Essink-Bot M, Kruijshaar M and Bac D. Different perceptions of the burden of upper GI endoscopy: An empirical study in three groups. Qual Life Res J. 2016; 16:1309–18.
- Feyzullah E, Toros A, Aydo G, Bektafi H and Ozcani O. Assesment of anxiety levels in patients during elective upper gastrointestinal endoscopy and colonoscopy, Turk J Gastroenterol. 2010; 21(1):29- 33.
- Gomez M and Llach J. Role of nurses in gastrointestinal endoscopy, Gastroenterol Hepatol J. 2009; 32(1):44-7.
- Hakan U, Gamze O and Haldun S. Departments of gastroenterology, Psychiatry and Internal Medicine J: Baflkent University School of Medicine, Ankara. 2012; 20(3): 67-71.
- Hashimoto T, Adachi K and Ishimura N. Safety and efficacy of glucagon as a pre medication for upper gastrointestinal endoscopy: A comparative study with butyl scopolamine bromide, Aliment Pharmacol Ther J. 2002; 16(1): 111-8.
- Hiremath P, Mohite V and Pawar S. A study to assess the knowledge, attitude and pre procedure anxiety level of patient undergoing upper gastrointestinal endoscopy, The American Journal of Gastroenterology.2016; 5(6):242-9.
- Jones M, Ebert C and Solan T .Relation between patient's anxiety and elective upper gastro endoscopy, Clinic Gastroenterol J. 2004; 38(1): 35-40.
- Kennedy J, Oconnor P and Sadovnick Upper gastrointestinal diseases influenced by place of residence, American Journal of Nursing. 2006; 93 (9): 162-7.
- Kiyohara Y and Kayanolk M. Anxiety in preoperative anesthetic endoscopy procedure, Rev Hosp Clinic Med Sao Paulo J. 2004; 11 (2):51-6.
- Koloski N and Boyce P. Epidemiology and health care seeking in the functional GI disorders: A population based study, Gastroenterol J. 2009; 97: 2290-9.
- Lee D, Chan A and Wong S. Can visual distraction decrease the dose of patient-controlled sedation required during endoscopy? A prospective randomized controlled trial, Endoscopy J. 2004; 36: 197-201.
- Lee Y, Anhn M, Kim E, Kweon H and Yoon M. The effect of preparatory education program on discomfort and retching of examinees during upper gastrointestinal endoscopy, Korean J Fam Med. 2012; 33(4): 219-28.
- Mohamad H, Mohamad Z and Sayed Z. Impact of a designed nursing teaching protocol on nurse's performance and patient's outcome in Esophagogastroduodenoscopy, Journal of American Science. 2014; 10 (10):56-65.
- Mohamed S. Effectiveness of structured teaching program on knowledge, anxiety state and tolerance for patients with gastrointestinal endoscopy, Life Science J. 2016; 13(9):9-17. Available

at: www.lifesciencesite.com.

- Mulcahy H and Kelly P. Factors associated with tolerance to, and discomfort with unsedated diagnostic gastroscopy, Scand J Gastroentrol. 2010; 36:1352-7.
- Muzzarelli L, Force M and Sebold M. Aromatherapy and reducing pre- gastrointestinal endoscopy anxiety: A controlled prospective study, Gastroenterol Nurs J. 2006; 29: 466-71.
- Pehlivan S, Ovayolu N and Koruk M. Effect of providing information to patient about upper gastrointestinal endoscopy on the patient's perception, compliance and anxiety level associated with the procedure, Journal of Clinical Nursing. 2011; 22(1):10-27.
- Peter B. Cotton N, Christopher B, Robert H and Brian P. The Fundamental of upper GI endoscopy, Philadelphia: Lippincott Williams
- Prabhuswami H, Vaishali R and Prakash N. Knowledge, attitude and pre-procedure anxiety level of patient undergoing upper gastrointestinal endoscopy in Krishna Hospital, India End J. 2016; 5 (6):243-9.
- Previti G, Bianchini O, Virzi A, Petralia A and Aguglia E. Identifying people at risk for anxiety in patients undergoing endoscopic procedures, Ann Depress Anxiety J. 2016; 3(1):1072-5.
- Quick J. Social support, secure attachments and health. Handbook, Philadelphia, Lippincott. 2006; 2(3): 169-72.
- Rohde H. Anxiety before gastrointestinal endoscopy is a significant problem, Int End J. 2011; 115(2):1539-44.
- Rudin D. Frequently overlooked and rarely listened to: Music therapy in gastrointestinal endoscopic procedures, World J Gastroenterol. 2007; 13(33): 4518- 33.
- Ruhl E and Everhart J. Assess indications and outcomes needs of GI endoscopy. 2016 3(1):184-6.Availableat: http://www3.niddk.nih.
- Salvetto M. The nurse's role in endoscopic ultra sonograph, Minerva Med J.2007;98(1):269-75.
- Seda P, Nimet O, Mehmet K, Yavuz P and Murat T. Effect of providing information to the patient about upper gastrointestinal endoscopy on the patient's perception, compliance and anxiety level associated with the procedure, Turk J Gastroenter. 2011; 22 (1):10-7.
- Sewitch M, Gong S, Barkun A, Hilsden R and Dube C. Literature review of quality in upper and 1ower gastrointestinal endoscopy from the patient perspective, Can J Gastroenterol. 2011; 25(1):681- 5.
- Spielberger C, Gorsuch R and Lushene R. Handbook for the IPAT anxiety scale. 2nd ed. Institute for personality and ability testing. 1970.
- Thomas S and Sugirtha V. Effectiveness of structured teaching programme on knowledge and anxiety of patients undergoing endoscopy at a gastroenterology centre of a tertiary care hospital, International Journal of Science and Research. 2015; 4(6): 1017-20. Available at: www.ijsr.net.
- Tina D and Steven P. An evaluation of a teaching intervention to improve the practice, Journal of Clinical Nursing. 2006; 10(5):682- 96.
- Trevisani L, Sartori S and Putinati S. Assessment of anxiety level in patients during diagnostic endoscopy. Recenti Prog Med J. 2002; 93(1): 240-4.
- Van L, Rossum G and Oijen M. Anxiety, depression and psychotropic medication use in patients with persistent upper gastrointestinal symptoms, Aliment Pharma TherJ. 2005; 21:1001-6.
- Vliet M, Grypdonck M and Zuuren F. Preparing patients for gastrointestinal endoscopy: The

influence of information in medical situations, Patient Educ Couns J. 2004; 52: 23-30.

- Winslow E. Patient education materials in upper GI endoscopy, American J of Nursing. 2004; 102 (17): 33-8.
- Yoenlee J, Anhn M, Kim D and Kweon H. The effect of preparatory education program on discomfort and retching of examinees during upper gastrointestinal endoscopy, Korean J Fam Med. 2012; 33(4): 219-28. Available at: http://synapse.koreamed.org.